

Quantum Processes and Computation

Assignment 1 – MT 2026

Unless otherwise noted, the following exercises are taken from *Picturing Quantum Software* (see github.com/zxcalc/book). There might be slight differences between the exercises on this problem sheet and how they appear in the book. **In that case, complete the version from this problem sheet and not the book version.** For some exercises, it may be useful to locate the exercise in the book for additional results and explanation.




Exercise 2.2. Write $\cos \alpha$ and $\sin \alpha$ as a sum of complex phases with rational coefficients.

Exercise 2.4. Let $M : H \rightarrow H$ be a linear map such that $M^\dagger M = MM^\dagger$. Then $M = \sum_j \lambda_j |\phi_j\rangle\langle\phi_j|$ for some set of complex numbers $\{\lambda_j\}_j$ and ONB $\{|\phi_j\rangle\}_j$. Show that:

- a) M is self-adjoint iff all $\lambda_j \in \mathbb{R}$
- b) M is positive iff all $\lambda_j \in \mathbb{R}_{\geq 0}$
- c) M is a projector iff all $\lambda_j \in \{0, 1\}$
- d) M is a unitary iff all $\lambda_j \in U(1)$

where $\mathbb{R}_{\geq 0}$ is the set of all real numbers ≥ 0 and $U(1) := \{e^{i\alpha} \mid \alpha \in [0, 2\pi)\}$ is the set of all **phases**.



$$\Phi_{abc}^{de} = \sum_{xyz} f_{ab}^{xd} g_{xy}^{ez} h_{cz}^y \quad (2.3)$$

Exercise 2.6. Write an expression for (2.2) using $\otimes$, $\circ$, SWAP, and $\text{Tr}(-)$.

Exercise 2.7. Show that for a normalised state $|\psi\rangle$, the operator $|\psi\rangle\langle\psi|$ is a projector with image $S := \{\lambda|\psi\rangle \mid \lambda \in \mathbb{C}\}$.

Exercise 2.8. Show that, for the following antipodal states on the Bloch sphere:

$$|\phi_0\rangle = \cos \frac{\theta}{2} |0\rangle + \sin \frac{\theta}{2} e^{i\alpha} |1\rangle$$

$$|\phi_1\rangle = \cos \frac{\theta + \pi}{2} |0\rangle + \sin \frac{\theta + \pi}{2} e^{i\alpha} |1\rangle$$

we have $\langle \phi_0 | \phi_1 \rangle = 0$.

Exercise 2.9. The standard basis $|0\rangle, |1\rangle$ corresponds to the Z axis of the Bloch sphere. Show that the following states


$$|+\rangle := \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$$

$$|-\rangle := \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$$

$$|+i\rangle := \frac{1}{\sqrt{2}}(|0\rangle + i|1\rangle)$$

$$|-i\rangle := \frac{1}{\sqrt{2}}(|0\rangle - i|1\rangle)$$

correspond to the X and Y axes, respectively. That is, show they are located on the Bloch sphere as follows:



Exercise 2.12. Define the following 3 measurements on a pair of qubits:

$$\mathcal{A} = \left\{ \begin{array}{c} \text{---} \triangleleft_i \triangleleft_i \text{---} \\ \text{---} \end{array} \right\}_i$$

$$\mathcal{B} = \left\{ \begin{array}{c} \text{---} \\ \text{---} \triangleleft_j \triangleleft_j \text{---} \end{array} \right\}_j$$

$$\mathcal{C} = \left\{ \begin{array}{c} \text{---} \triangleleft_i \triangleleft_i \text{---} \\ \text{---} \triangleleft_j \triangleleft_j \text{---} \end{array} \right\}_{ij}$$

Show that the probability of getting outcome i for $\mathcal{A}$ then getting outcome j when measuring $\mathcal{B}$ on the resulting state is the same as the probability of getting outcome (i, j) for $\mathcal{C}$.