

LECTURE 7: PROPOSITIONAL LOGIC (1)

Software Engineering
Mike Wooldridge

- The *syntax* of a logic defines the syntactically acceptable objects of the language, which are properly called *well-formed formulae* (wff). (We shall just call them formulae.)
- The *semantics* of a logic associate each formula with a *meaning*.
- The *proof theory* is concerned with manipulating formulae according to certain rules.

1 What is a Logic?

- When most people say 'logic', they mean either *propositional logic* or *first-order predicate logic*.
- However, the precise definition is quite broad, and literally hundreds of logics have been studied by philosophers, computer scientists and mathematicians.
- Any 'formal system' can be considered a logic if it has:
 - a well-defined *syntax*;
 - a well-defined *semantics*; and
 - a well-defined *proof-theory*.

2 Propositional Logic

- The simplest, and most abstract logic we can study is called *propositional logic*.
- **Definition:** A *proposition* is a statement that can be either *true* or *false*; it must be one or the other, and it cannot be both.
- **EXAMPLES.** The following are propositions:
 - the reactor is on;
 - the wing-flaps are up;
 - John Major is prime minister.whereas the following are not:
 - are you going out somewhere?
 - $2+3$

- It is possible to determine whether any given statement is a proposition by prefixing it with:

It is true that ...

and seeing whether the result makes grammatical sense.

- We now define *atomic* propositions. Intuitively, these are the set of smallest propositions.
- **Definition:** An *atomic proposition* is one whose truth or falsity does not depend on the truth or falsity of any other proposition.
- So all the above propositions are atomic.

2.1 The Connectives

- Now, the study of atomic propositions is pretty boring. We therefore now introduce a number of *connectives* which will allow us to build up *complex propositions*.
- The connectives we introduce are:
 - \wedge and (& or .)
 - \vee or (| or +)
 - \neg not (\sim)
 - \Rightarrow implies (\supset or \rightarrow)
 - \Leftrightarrow iff
- (Some books use other notations; these are given in parentheses.)

- Now, rather than write out propositions in full, we will abbreviate them by using *propositional variables*.

- It is standard practice to use the lower-case roman letters

p, q, r, \dots

to stand for propositions.

- If we do this, we must define what we mean by writing something like:

Let p be *John Major is prime Minister*.

- Another alternative is to write something like *reactor_is_on*, so that the interpretation of the propositional variable becomes obvious.

And

- Any two propositions can be combined to form a third proposition called the *conjunction* of the original propositions.
- **Definition:** If p and q are arbitrary propositions, then the *conjunction* of p and q is written

$$p \wedge q$$

and will be true iff both p and q are true.

- We can summarise the operation of \wedge in a *truth table*. The idea of a truth table for some formula is that it describes the behaviour of a formula under all possible interpretations of the primitive propositions the are included in the formula.
- If there are n different atomic propositions in some formula, then there are 2^n different lines in the truth table for that formula. (This is because each proposition can take one 1 of 2 values — *true* or *false*.)
- Let us write T for truth, and F for falsity. Then the truth table for $p \wedge q$ is:

p	q	$p \wedge q$
F	F	F
F	T	F
T	F	F
T	T	T

- The operation of \vee is summarised in the following truth table:

p	q	$p \vee q$
F	F	F
F	T	T
T	F	T
T	T	T

Or

- Any two propositions can be combined by the word 'or' to form a third proposition called the *disjunction* of the originals.
- **Definition:** If p and q are arbitrary propositions, then the *disjunction* of p and q is written

$$p \vee q$$

and will be true iff either p is true, or q is true, or both p and q are true.

If... Then...

- Many statements, particularly in mathematics, are of the form:

if p is true then q is true.

Another way of saying the same thing is to write:

p implies q .

- In propositional logic, we have a connective that combines two propositions into a new proposition called the *conditional*, or *implication* of the originals, that attempts to capture the sense of such a statement.

- **Definition:** If p and q are arbitrary propositions, then the *conditional* of p and q is written

$$p \Rightarrow q$$

and will be true iff either p is false or q is true.

- The truth table for \Rightarrow is:

p	q	$p \Rightarrow q$
F	F	T
F	T	T
T	F	F
T	T	T

Iff

- Another common form of statement in maths is:

p is true if, and only if, q is true.

- The sense of such statements is captured using the *biconditional* operator.
- **Definition:** If p and q are arbitrary propositions, then the *biconditional* of p and q is written:

$$p \Leftrightarrow q$$

and will be true iff either:

1. p and q are both true; or
2. p and q are both false.

- The \Rightarrow operator is the hardest to understand of the operators we have considered so far, and yet it is extremely important.
- If you find it difficult to understand, just remember that the $p \Rightarrow q$ means 'if p is true, then q is true'.
If p is false, then we don't care about q , and by default, make $p \Rightarrow q$ evaluate to T in this case.
- Terminology: if ϕ is the formula $p \Rightarrow q$, then p is the *antecedent* of ϕ and q is the *consequent*.

- The truth table for \Leftrightarrow is:

p	q	$p \Leftrightarrow q$
F	F	T
F	T	F
T	F	F
T	T	T

- If $p \Leftrightarrow q$ is true, then p and q are said to be *logically equivalent*. They will be true under exactly the same circumstances.

Not

- All of the connectives we have considered so far have been *binary*: they have taken *two* arguments.
- The final connective we consider here is *unary*. It only takes *one* argument.
- Any proposition can be prefixed by the word 'not' to form a second proposition called the *negation* of the original.
- **Definition:** If p is an arbitrary proposition then the *negation* of p is written

$$\neg p$$

and will be true iff p is false.

- Truth table for \neg :

p	$\neg p$
F	T
T	F

3 Tautologies & Consistency

- Given a particular formula, can you tell if it is true or not?
- No — you usually need to know the truth values of the component atomic propositions in order to be able to tell whether a formula is true.
- **Definition:** A *valuation* is a function which assigns a truth value to each primitive proposition.
- In Modula-2, we might write:


```
PROCEDURE Val(p : AtomicProp) :
  BOOLEAN;
```
- Given a valuation, we can say for any formula whether it is true or false.

Comments

- We can *nest* complex formulae as deeply as we want.
- We can use *parentheses* i.e., $()$, to *disambiguate* formulae.
- **EXAMPLES.** If p, q, r, s and t are atomic propositions, then all of the following are formulae:

- $p \wedge q \Rightarrow r$
- $p \wedge (q \Rightarrow r)$
- $(p \wedge (q \Rightarrow r)) \vee s$
- $((p \wedge (q \Rightarrow r)) \vee s) \wedge t$

whereas none of the following is:

- $p \wedge$
- $p \wedge q)$
- $p \neg$

- **EXAMPLE.** Suppose we have a valuation v , such that:

$$\begin{aligned} v(p) &= F \\ v(q) &= T \\ v(r) &= F \end{aligned}$$

Then we truth value of $(p \vee q) \Rightarrow r$ is evaluated by:

$$\begin{aligned} (v(p) \vee v(q)) \Rightarrow v(r) & \quad (1) \\ = (F \vee T) \Rightarrow F & \quad (2) \\ = T \Rightarrow F & \quad (3) \\ = F & \quad (4) \end{aligned}$$

Line (3) is justified since we know that $F \vee T = T$.

Line (4) is justified since $T \Rightarrow F = F$.

If you can't see this, look at the truth tables for \vee and \Rightarrow .

- When we consider formulae in terms of interpretations, it turns out that some have interesting properties.
- **Definition:**
 1. A formula is a *tautology* iff it is true under *every* valuation;
 2. A formula is *consistent* iff it is true under *at least one* valuation;
 3. A formula is *inconsistent* iff it is not made true under *any* valuation.
- Now, each line in the truth table of a formula corresponds to a valuation.
- So, we can use truth tables to determine whether or not formulae are tautologies.