

Full Abstraction for Nominal General References

Nikos Tzevelekos

`nikt@comlab.ox.ac.uk`

LICS, Wroclaw, July 2007

Full Abstraction for Nominal General References – Overview

This talk is about formulating a fully-abstract semantics of nominal general references using nominal games.

Full Abstraction for Nominal General References – Overview

This talk is about formulating a fully-abstract semantics of nominal general references using nominal games.

We will be talking about:

- Nominal Sets (Gabbay, Pitts)
- A functional higher-order language with nominal general references (Pitts, Stark, NT), the $\nu\rho$ -calculus:
 - *good variables (references)*,
 - *reference-equality tests*.
- Nominal Games (Abramsky, Ghica, Murawski, Ong, Stark, NT)

Nominal Sets

Assume a countable set of types TY and for each $A \in TY$ a countably infinite set of *names* N_A . Take,

- $N \triangleq \bigsqcup_{A \in TY} N_A$ to be the set of *general names*,
- and $PERM(N) = \bigoplus_{A \in TY} PERM(N_A)$.

Nominal Sets

Assume a countable set of types TY and for each $A \in \text{TY}$ a countably infinite set of *names* N_A . Take,

- $\mathbf{N} \triangleq \bigsqcup_{A \in \text{TY}} N_A$ to be the set of *general names*,
- and $\text{PERM}(\mathbf{N}) = \bigoplus_{A \in \text{TY}} \text{PERM}(N_A)$.

A *nominal set* X is a set equipped with an action from $\text{PERM}(\mathbf{N})$,

$$_ \circ _ : \text{PERM}(\mathbf{N}) \times X \rightarrow X \quad (\text{e.g. } \pi \circ x)$$

such that all $x \in X$ have *finite strong support* $\mathbf{S}(x) \subseteq \mathbf{N}$,

$$\forall \pi \in \text{PERM}(\mathbf{N}). \pi \circ x = x \iff \forall \alpha \in \mathbf{S}(x). \pi(\alpha) = \alpha$$

Nominal Sets

Assume a countable set of types TY and for each $A \in \text{TY}$ a countably infinite set of *names* N_A . Take,

- $N \triangleq \bigsqcup_{A \in \text{TY}} N_A$ to be the set of *general names*,
- and $\text{PERM}(N) = \bigoplus_{A \in \text{TY}} \text{PERM}(N_A)$.

A *nominal set* X is a set equipped with an action from $\text{PERM}(N)$,

$$_ \circ _ : \text{PERM}(N) \times X \rightarrow X \quad (\text{e.g. } \pi \circ x)$$

such that all $x \in X$ have *finite strong support* $S(x) \subseteq N$,

$$\forall \pi \in \text{PERM}(N). \pi \circ x = x \iff \forall \alpha \in S(x). \pi(\alpha) = \alpha$$

$\rightsquigarrow N^\#$, the set of *finite lists of distinct names*, is a nominal set.
(denote names by α, β , etc. and lists of names by $\vec{\alpha}, \vec{\beta}$, etc.)

\rightsquigarrow If X, Y nominal sets then $X \times Y$ a nominal set.

The $\nu\rho$ -calculus –use names for general references!

The $\nu\rho$ -calculus is a functional calculus with nominal references.

$$\mathbf{TY} \ni A, B ::= \mathbb{1} \mid \mathbb{N} \mid [A] \mid A \rightarrow B \mid A \otimes B$$

The $\nu\rho$ -calculus –use names for general references!

The $\nu\rho$ -calculus is a functional calculus with nominal references.

TY $\ni A, B ::= \mathbb{1} \mid \mathbb{N} \mid [A] \mid A \rightarrow B \mid A \otimes B$

TE $\ni M, N ::= x \mid \lambda x.M \mid M N$ λ -term
| skip return
| $\tilde{n} \mid \text{pred } M \mid \text{succ } N$ arithmetic
| if0 M then N_1 else N_2 if_then_else
| $\langle M, N \rangle \mid \text{fst } M \mid \text{snd } N$ pair / projections

The $\nu\rho$ -calculus –use names for general references!

The $\nu\rho$ -calculus is a functional calculus with nominal references.

$$\mathbf{TY} \ni A, B ::= \mathbb{1} \mid \mathbb{N} \mid [A] \mid A \rightarrow B \mid A \otimes B$$

$\mathbf{TE} \ni M, N ::= x \mid \lambda x.M \mid M N$	λ -term
$\mid \text{skip}$	return
$\mid \tilde{n} \mid \text{pred } M \mid \text{succ } N$	arithmetic
$\mid \text{if0 } M \text{ then } N_1 \text{ else } N_2$	if_then_else
$\mid \langle M, N \rangle \mid \text{fst } M \mid \text{snd } N$	pair / projections
$\mid \alpha$	name, $\alpha \in \mathbb{N} = \bigsqcup_{A \in \mathbf{TY}} \mathbb{N}_A$
$\mid \nu\alpha.M$	ν -abstraction
$\mid [M = N]$	name-equality test
$\mid M := N$	update
$\mid !M$	dereferencing

The $\nu\rho$ -calculus –use names for general references!

The $\nu\rho$ -calculus is a functional calculus with nominal references.

$$\mathbf{TY} \ni A, B ::= \mathbb{1} \mid \mathbb{N} \mid [A] \mid A \rightarrow B \mid A \otimes B$$

$\mathbf{TE} \ni M, N ::=$	$x \mid \lambda x.M \mid M N$	λ -term
	$\mid \text{skip}$	return
	$\mid \tilde{n} \mid \text{pred } M \mid \text{succ } N$	arithmetic
	$\mid \text{if0 } M \text{ then } N_1 \text{ else } N_2$	if_then_else
	$\mid \langle M, N \rangle \mid \text{fst } M \mid \text{snd } N$	pair / projections
	$\mid \alpha$	name, $\alpha \in \mathbb{N} = \bigsqcup_{A \in \mathbf{TY}} \mathbb{N}_A$
	$\mid \nu\alpha.M$	ν -abstraction
	$\mid [M = N]$	name-equality test
	$\mid M := N \mid !M$	update / dereferencing

$$\mathbf{VA} \ni V, W ::= \tilde{n} \mid \text{skip} \mid \alpha \mid x \mid \lambda x.M \mid \langle V, W \rangle$$

The $\nu\rho$ -calculus: Typed Terms

Terms are typed in environments $(\vec{\alpha}, \Gamma)$ consisting of:

- a *list* $\vec{\alpha}$ of distinct names ($\vec{\alpha} \in \mathbf{N}^\#$)
- a set Γ of variable-type pairs

$\vec{\alpha} \mid \Gamma \vdash M : A$

\rightsquigarrow free vars in Γ

\rightsquigarrow (free) names in $\vec{\alpha}$

The $\nu\rho$ -calculus: Typed Terms

Terms are typed in environments $(\vec{\alpha}, \Gamma)$ consisting of:

- a *list* $\vec{\alpha}$ of distinct names ($\vec{\alpha} \in \mathbf{N}^\#$)
- a set Γ of variable-type pairs

$\vec{\alpha} \mid \Gamma \vdash M : A$
 \rightsquigarrow free vars in Γ
 \rightsquigarrow (free) names in $\vec{\alpha}$

$$\frac{}{\vec{\alpha} \mid \Gamma, x : A \vdash x : A}$$

$$\frac{}{\vec{\alpha} \mid \Gamma \vdash \alpha : [A]} \quad \begin{matrix} \alpha \in \mathcal{S}(\vec{\alpha}) \\ \wedge \alpha \in \mathbf{N}_A \end{matrix}$$

$$\frac{\vec{\alpha}\alpha \mid \Gamma \vdash M : B}{\vec{\alpha} \mid \Gamma \vdash \nu\alpha.M : B}$$

$$\frac{\vec{\alpha} \mid \Gamma \vdash M : [A] \quad \vec{\alpha} \mid \Gamma \vdash N : [A]}{\vec{\alpha} \mid \Gamma \vdash [M = N] : \mathbb{N}}$$

$$\frac{\vec{\alpha} \mid \Gamma \vdash M : [A]}{\vec{\alpha} \mid \Gamma \vdash !M : A}$$

$$\frac{\vec{\alpha} \mid \Gamma \vdash M : [A] \quad \vec{\alpha} \mid \Gamma \vdash N : A}{\vec{\alpha} \mid \Gamma \vdash M := N : \mathbb{1}}$$

The $\nu\rho$ -calculus: Reduction

The reduction calculus is defined in store environments S :

$$S ::= \epsilon \mid \alpha, S \mid \alpha :: V, S$$

with their domains being lists of distinct names.

$$\text{EQ} \frac{}{S \vDash [\alpha = \beta]} \longrightarrow S \vDash \tilde{n} \quad \begin{array}{l} n=1 \text{ if } \alpha \neq \beta \\ n=0 \text{ if } \alpha = \beta \end{array}$$

$$\text{NEW} \frac{}{S \vDash \nu\alpha.M} \longrightarrow S, \beta \vDash (\alpha \beta) \circ M \quad \beta \notin \mathbf{S}(S)$$

$$\text{DRF} \frac{}{S, \alpha :: V, S' \vDash !\alpha} \longrightarrow S, \alpha :: V, S' \vDash V$$

$$\text{UPD} \frac{}{S, \alpha(:: W), S' \vDash \alpha := V} \longrightarrow S, \alpha :: V, S' \vDash \text{skip}$$

$$\text{LAM} \frac{}{S \vDash (\lambda x.M) V} \longrightarrow S \vDash M\{V/x\}$$

The $\nu\rho$ -calculus: Reduction

The reduction calculus is defined in store environments S :

$$S ::= \epsilon \mid \alpha, S \mid \alpha :: V, S$$

with their domains being lists of distinct names.

$$\text{EQ} \frac{}{S \vDash [\alpha = \beta]} \longrightarrow S \vDash \tilde{n} \quad \begin{array}{l} n=1 \text{ if } \alpha \neq \beta \\ n=0 \text{ if } \alpha = \beta \end{array}$$

$$\text{NEW} \frac{}{S \vDash \nu\alpha.M} \longrightarrow S, \beta \vDash (\alpha \beta) \circ M \quad \beta \notin \mathbf{s}(S)$$

$$\text{DRF} \frac{}{S, \alpha :: V, S' \vDash !\alpha} \longrightarrow S, \alpha :: V, S' \vDash V$$

$$\text{UPD} \frac{}{S, \alpha(:: W), S' \vDash \alpha := V} \longrightarrow S, \alpha :: V, S' \vDash \text{skip}$$

$$\text{LAM} \frac{}{S \vDash (\lambda x.M) V} \longrightarrow S \vDash M\{V/x\}$$

$\nu\rho$ -calculus : Observational Equivalence

Reduction yields the following notion of equivalence.

For typed terms $\vec{\alpha} \mid \Gamma \vdash M : A$ and $\vec{\alpha} \mid \Gamma \vdash N : A$,

$$\vec{\alpha} \mid \Gamma \vdash M \approx N \iff$$

$$\forall C[_] : \mathbb{N}. (\exists S'. \vdash C[M] \longrightarrow S' \vdash \tilde{0})$$

$$\implies (\exists S''. \vdash C[N] \longrightarrow S'' \vdash \tilde{0})$$

where $C[_]$ is a variable- and name-closing context.

$\nu\rho$ -calculus : Observational Equivalence

Reduction yields the following notion of equivalence.

For typed terms $\vec{\alpha} \mid \Gamma \vdash M : A$ and $\vec{\alpha} \mid \Gamma \vdash N : A$,

$$\vec{\alpha} \mid \Gamma \vdash M \approx N \iff$$

$$\forall C[-] : \mathbb{N}. (\exists S'. \vdash C[M] \longrightarrow S' \vdash \tilde{0})$$

$$\implies (\exists S''. \vdash C[N] \longrightarrow S'' \vdash \tilde{0})$$

where $C[-]$ is a variable- and name-closing context.

For example,

$$\nu\alpha.M \cong M \quad \text{if } \alpha \notin \mathcal{S}(M)$$

$$\nu\alpha, \beta.M \cong \nu\beta, \alpha.M$$

Nominal Games: Arenas and Prearenas

Game semantics gained prominence in the mid-90's by providing the first fully abstract semantics for PCF. Since then, game semantics have been used in order to model fully-abstractly languages with a wide range of computational effects –mostly in LICS.

Nominal Games: Arenas and Prearenas

A (nominal) game can be described by *plays* –sequences of moves played in alternation by *Opponent* and *Player*– on a *prearena*.

An *arena* $A \triangleq (M_A, I_A, \vdash_A, \lambda_A)$ is given by:

Nominal Games: Arenas and Prearenas

A (nominal) game can be described by *plays* –sequences of moves played in alternation by *Opponent* and *Player*– on a *prearena*.

An *arena* $A \triangleq (M_A, I_A, \vdash_A, \lambda_A)$ is given by:

- A nominal set M_A of moves,
- A nominal subset $I_A \subseteq M_A$ of *initial* moves,
- A nominal *justification* relation $\vdash_A \subseteq M_A \times (M_A \setminus I_A)$,
- A nominal *labeling* function $\lambda_A : M_A \rightarrow \{O, P\} \times \{A, Q\}$.

Nominal Games: Arenas and Prearenas

A (nominal) game can be described by *plays* –sequences of moves played in alternation by *Opponent* and *Player*– on a *prearena*.

An *arena* $A \triangleq (M_A, I_A, \vdash_A, \lambda_A)$ is given by:

- A nominal set M_A of moves,
- A nominal subset $I_A \subseteq M_A$ of *initial* moves,
- A nominal *justification* relation $\vdash_A \subseteq M_A \times (M_A \setminus I_A)$,
- A nominal *labeling* function $\lambda_A : M_A \rightarrow \{O, P\} \times \{A, Q\}$.

↪ For each $m \in (M_A \setminus I_A)$: $i_A \vdash_A m_1 \vdash_A \cdots \vdash_A m_i \vdash_A m$

↪ Initial moves are *P*-Answers, and whenever $m_1 \vdash_A m_2$,

m_1, m_2 by different players , m_1 Answer $\implies m_2$ Question

Nominal Games: Arenas and Prearenas

A (nominal) game can be described by *plays* –sequences of moves played in alternation by *Opponent* and *Player*– on a *prearena*.

An *arena* $A \triangleq (M_A, I_A, \vdash_A, \lambda_A)$ is given by:

- A nominal set M_A of moves,
- A nominal subset $I_A \subseteq M_A$ of *initial* moves,
- A nominal *justification* relation $\vdash_A \subseteq M_A \times (M_A \setminus I_A)$,
- A nominal *labeling* function $\lambda_A : M_A \rightarrow \{O, P\} \times \{A, Q\}$.

↪ For each $m \in (M_A \setminus I_A)$: $i_A \vdash_A m_1 \vdash_A \cdots \vdash_A m_i \vdash_A m$

↪ Initial moves are *O-Questions*, and whenever $m_1 \vdash_A m_2$,

m_1, m_2 by different players , m_1 Answer $\implies m_2$ Question

Basic Arenas, Prearenas

$$\frac{1}{M_1 \triangleq \{*\}}$$

$$I_1 \triangleq \{*\}$$

$$\frac{\mathbb{N}}{M_{\mathbb{N}} \triangleq \mathbb{N}}$$

$$I_{\mathbb{N}} \triangleq \mathbb{N}$$

$$\frac{N_A}{M_{N_A} \triangleq N_A}$$

$$I_{N_A} \triangleq N_A$$

$$\frac{\mathbb{N} \Rightarrow N_A}{*}$$

PA

n *α*

OQ

PA

Basic Arenas, Prearenas

$$\frac{1}{M_1 \triangleq \{*\}$$

$$I_1 \triangleq \{*\}$$




$$\frac{\mathbb{N}}{M_{\mathbb{N}} \triangleq \mathbb{N}}$$

$$I_{\mathbb{N}} \triangleq \mathbb{N}$$

$$\frac{N_A}{M_{N_A} \triangleq N_A}$$

$$I_{N_A} \triangleq N_A$$

$$\frac{\mathbb{N} \Rightarrow N_A}{\begin{array}{c} * \\ \curvearrowright \\ n \\ \curvearrowright \\ \alpha \end{array}} \quad \begin{array}{l} PA \\ OQ \\ PA \end{array}$$



Basic Arenas, Prearenas

$$\frac{1}{M_1 \triangleq \{*\}$$

$$I_1 \triangleq \{*\}$$







$$\frac{\mathbb{N}}{M_{\mathbb{N}} \triangleq \mathbb{N}}$$

$$I_{\mathbb{N}} \triangleq \mathbb{N}$$

$$\frac{N_A}{M_{N_A} \triangleq N_A}$$

$$I_{N_A} \triangleq N_A$$


$$\frac{\mathbb{N} \Rightarrow N_A}{\begin{array}{c} * \\ \curvearrowright \\ n \\ \curvearrowright \\ \alpha \end{array}} \quad \begin{array}{l} PA \\ OQ \\ PA \end{array}$$



OQ

P

Moves and Plays

A *move-with-names* of a prearena A is: $m_{\vec{\alpha}}$  $\vec{\alpha} \in \mathbb{N}^\#$ *-local state*
 $m \in M_A$

An $\vec{\alpha}$ -*play* on A is a sequence of moves-with-names such that:

- moves are OP -alternating,
- every move is justified by a previous one,
- except from the first move, which is initial and has local state $\vec{\alpha}$,
- Visibility and Well-Bracketing are satisfied.
- O -moves cannot change the local state,
- P -moves can add fresh names to the local state,
- P -moves must add fresh names to the local state to use them,

e.g. $N_A \xrightarrow{\beta} N_A \Rightarrow N_A$

β


OQ

PA

OQ

PA

Moves and Plays

A *move-with-names* of a prearena A is: $m_{\vec{\alpha}}$  $\vec{\alpha} \in \mathbb{N}^\#$ *-local state*
 $m \in M_A$

An $\vec{\alpha}$ -*play* on A is a sequence of moves-with-names such that:

- moves are OP -alternating,
- every move is justified by a previous one,
- except from the first move, which is initial and has local state $\vec{\alpha}$,
- Visibility and Well-Bracketing are satisfied.
- O -moves cannot change the local state,
- P -moves can add fresh names to the local state,
- P -moves must add fresh names to the local state to use them,

e.g.

$$N_A \longrightarrow N_A \Rightarrow N_A$$

β




OQ

PA

OQ

PA

Moves and Plays


A *move-with-names* of a prearena A is: $m_{\vec{\alpha}}$  $\vec{\alpha} \in \mathbb{N}^\#$ *-local state*
 $m \in M_A$

An $\vec{\alpha}$ -*play* on A is a sequence of moves-with-names such that:

- moves are OP -alternating,
- every move is justified by a previous one,
- except from the first move, which is initial and has local state $\vec{\alpha}$,
- Visibility and Well-Bracketing are satisfied.
- O -moves cannot change the local state,
- P -moves can add fresh names to the local state,
- P -moves must add fresh names to the local state to use them,

e.g.

$$N_A \longrightarrow N_A \Rightarrow N_A$$



OQ

PA

OQ

PA

Moves and Plays


A *move-with-names* of a prearena A is: $m_{\vec{\alpha}}$ $\vec{\alpha} \in \mathbb{N}^\#$ *-local state*
 $m \in M_A$

An $\vec{\alpha}$ -*play* on A is a sequence of moves-with-names such that:

- moves are OP -alternating,
- every move is justified by a previous one,
- except from the first move, which is initial and has local state $\vec{\alpha}$,
- Visibility and Well-Bracketing are satisfied.
- O -moves cannot change the local state,
- P -moves can add fresh names to the local state,
- P -moves must add fresh names to the local state to use them,

e.g.

$$N_A \longrightarrow N_A \Rightarrow N_A$$



Moves and Plays


A *move-with-names* of a prearena A is: $m_{\vec{\alpha}}$ $\vec{\alpha} \in \mathbb{N}^\#$ *-local state*
 $m \in M_A$

An $\vec{\alpha}$ -*play* on A is a sequence of moves-with-names such that:

- moves are OP -alternating,
- every move is justified by a previous one,
- except from the first move, which is initial and has local state $\vec{\alpha}$,
- Visibility and Well-Bracketing are satisfied.
- O -moves cannot change the local state,
- P -moves can add fresh names to the local state,
- P -moves must add fresh names to the local state to use them,

e.g.

$$N_A \longrightarrow N_A \Rightarrow N_A$$



Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

$$[s]_{\vec{\alpha}} \triangleq \{ \pi \circ s \mid \pi \circ \vec{\alpha} = \vec{\alpha} \}$$

Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

- If even-length $[s_1x_1]_{\vec{\alpha}}, [s_2x_2]_{\vec{\alpha}} \in \sigma$ and $[s_1]_{\vec{\alpha}} = [s_2]_{\vec{\alpha}}$ then $[s_1x_1]_{\vec{\alpha}} = [s_2x_2]_{\vec{\alpha}}$. (*determinacy*)

$$[s]_{\vec{\alpha}} \triangleq \{ \pi \circ s \mid \pi \circ \vec{\alpha} = \vec{\alpha} \}$$

Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

- If even-length $[s_1x_1]_{\vec{\alpha}}, [s_2x_2]_{\vec{\alpha}} \in \sigma$ and $[s_1]_{\vec{\alpha}} = [s_2]_{\vec{\alpha}}$ then $[s_1x_1]_{\vec{\alpha}} = [s_2x_2]_{\vec{\alpha}}$. (*determinacy*)
- *Innocence* and Totality.

$$[s]_{\vec{\alpha}} \triangleq \{ \pi \circ s \mid \pi \circ \vec{\alpha} = \vec{\alpha} \}$$

An $\vec{\alpha}$ -strategy σ on $A \rightarrow B$ is written $\sigma : A \rightarrow B$.

Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

- If even-length $[s_1x_1]_{\vec{\alpha}}, [s_2x_2]_{\vec{\alpha}} \in \sigma$ and $[s_1]_{\vec{\alpha}} = [s_2]_{\vec{\alpha}}$ then $[s_1x_1]_{\vec{\alpha}} = [s_2x_2]_{\vec{\alpha}}$. (*determinacy*)
- *Innocence* and Totality.

An $\vec{\alpha}$ -strategy σ on $A \rightarrow B$ is written $\sigma : A \rightarrow B$.

$$\left. \begin{array}{l} \sigma : A \rightarrow B \text{ an } \vec{\alpha}\text{-strategy} \\ \tau : B \rightarrow C \text{ an } \vec{\alpha}\text{-strategy} \end{array} \right\} \quad A \xrightarrow{\sigma} B \xrightarrow{\tau} C$$


Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

- If even-length $[s_1x_1]_{\vec{\alpha}}, [s_2x_2]_{\vec{\alpha}} \in \sigma$ and $[s_1]_{\vec{\alpha}} = [s_2]_{\vec{\alpha}}$ then $[s_1x_1]_{\vec{\alpha}} = [s_2x_2]_{\vec{\alpha}}$. (*determinacy*)
- *Innocence* and Totality.

An $\vec{\alpha}$ -strategy σ on $A \rightarrow B$ is written $\sigma : A \rightarrow B$.

$\left. \begin{array}{l} \sigma : A \rightarrow B \text{ an } \vec{\alpha}\text{-strategy} \\ \tau : B \rightarrow C \text{ an } \vec{\alpha}\text{-strategy} \end{array} \right\}$



Strategies

An $\vec{\alpha}$ -strategy σ is a prefix-closed and O-move-closed set of equivalence classes $[s]_{\vec{\alpha}}$ of $\vec{\alpha}$ -plays, satisfying:

- If even-length $[s_1x_1]_{\vec{\alpha}}, [s_2x_2]_{\vec{\alpha}} \in \sigma$ and $[s_1]_{\vec{\alpha}} = [s_2]_{\vec{\alpha}}$ then $[s_1x_1]_{\vec{\alpha}} = [s_2x_2]_{\vec{\alpha}}$. (*determinacy*)
- *Innocence* and Totality.

An $\vec{\alpha}$ -strategy σ on $A \rightarrow B$ is written $\sigma : A \rightarrow B$.



$\sigma; \tau : A \rightarrow C$ an $\vec{\alpha}$ -strategy, obtained by:

\rightsquigarrow *composing and hiding B-moves*

\rightsquigarrow *respecting Name Conditions*

Let $\mathcal{V}^{\vec{\alpha}}$ be the category of nominal arenas and $\vec{\alpha}$ -strategies

Translate types to arenas:

$$\llbracket 1 \rrbracket = 1 \quad \llbracket N \rrbracket = N \quad \llbracket [A] \rrbracket = N_A \quad \llbracket A \otimes B \rrbracket = \llbracket A \rrbracket \otimes \llbracket B \rrbracket$$

Let $\mathcal{V}^{\vec{\alpha}}$ be the category of nominal arenas and $\vec{\alpha}$ -strategies

Translate types to arenas:

$$[[1]] = 1 \quad [[N]] = N \quad [[A]] = N_A \quad [[A \otimes B]] = [[A]] \otimes [[B]]$$

$$[[A \rightarrow B]] = [[A]] \otimes \xi \Rightarrow [[B]] \otimes \xi \quad \xi = \bigotimes_A (N_A \Rightarrow [[A]])$$

– *solved as a recursive domain equation.*

Let $\mathcal{V}^{\vec{\alpha}}$ be the category of nominal arenas and $\vec{\alpha}$ -strategies

Translate types to arenas:

$$\llbracket \mathbf{1} \rrbracket = 1 \quad \llbracket \mathbf{N} \rrbracket = \mathbf{N} \quad \llbracket [A] \rrbracket = N_A \quad \llbracket A \otimes B \rrbracket = \llbracket A \rrbracket \otimes \llbracket B \rrbracket$$

$$\llbracket A \rightarrow B \rrbracket = \llbracket A \rrbracket \overset{\sim}{\Rightarrow} (\xi \Rightarrow \llbracket B \rrbracket \otimes \xi) \quad \xi = \bigotimes_A (N_A \Rightarrow \llbracket A \rrbracket)$$

We obtain a *store monad* T , taking $TA \triangleq \xi \Rightarrow (A \otimes \xi)$.

Semantics in $\mathcal{V}^{\vec{\alpha}}$


Let $\mathcal{V}^{\vec{\alpha}}$ be the category of nominal arenas and $\vec{\alpha}$ -strategies

Translate types to arenas:

$$\llbracket \mathbf{1} \rrbracket = 1 \quad \llbracket \mathbf{N} \rrbracket = \mathbf{N} \quad \llbracket [A] \rrbracket = N_A \quad \llbracket A \otimes B \rrbracket = \llbracket A \rrbracket \otimes \llbracket B \rrbracket$$

$$\llbracket A \rightarrow B \rrbracket = \llbracket A \rrbracket \overset{\sim}{\Rightarrow} (\xi \Rightarrow \llbracket B \rrbracket \otimes \xi) \quad \xi = \bigotimes_A (N_A \Rightarrow \llbracket A \rrbracket)$$

We obtain a *store monad* T , taking $TA \triangleq \xi \Rightarrow (A \otimes \xi)$.



Semantics in $\mathcal{V}^{\vec{\alpha}}$


Let $\mathcal{V}^{\vec{\alpha}}$ be the category of nominal arenas and $\vec{\alpha}$ -strategies

Translate types to arenas:

$$\llbracket \mathbf{1} \rrbracket = 1 \quad \llbracket \mathbf{N} \rrbracket = \mathbf{N} \quad \llbracket [A] \rrbracket = N_A \quad \llbracket A \otimes B \rrbracket = \llbracket A \rrbracket \otimes \llbracket B \rrbracket$$

$$\llbracket A \rightarrow B \rrbracket = \llbracket A \rrbracket \overset{\sim}{\Rightarrow} (\xi \Rightarrow \llbracket B \rrbracket \otimes \xi) \quad \xi = \bigotimes_A (N_A \Rightarrow \llbracket A \rrbracket)$$

We obtain a *store monad* T , taking $TA \triangleq \xi \Rightarrow (A \otimes \xi)$.



Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$$\frac{[[\alpha \mid x : A \vdash [x = \alpha] : \mathbb{N}]]}{}$$

$$N_A \longrightarrow T\mathbb{N}$$

$$\alpha^\alpha \quad (\xi \Rightarrow \mathbb{N} \otimes \xi)$$

$$\frac{[[\mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N}]]}{}$$

$$N_A \longrightarrow T\mathbb{N}$$

$$(\xi \Rightarrow \mathbb{N} \otimes \xi)$$

Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow T\mathbb{N}$

α^α $(\xi \Rightarrow \mathbb{N} \otimes \xi)$

 $*^\alpha$

$\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket$


$N_A \longrightarrow T\mathbb{N}$

$(\xi \Rightarrow \mathbb{N} \otimes \xi)$

Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow TN$




$\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow TN$

$(\xi \Rightarrow \mathbb{N} \otimes \xi)$

Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$$\frac{\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket}{N_A \longrightarrow TN}$$



$$\frac{\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket}{N_A \longrightarrow TN}$$




Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket$

$\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow TN$

$N_A \longrightarrow TN$



$(\xi \Rightarrow \mathbb{N} \otimes \xi)$



Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$

$\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket$

$\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow T\mathbb{N}$


$N_A \longrightarrow T\mathbb{N}$



Examples of semantics in the model $\mathcal{V} \triangleq \langle \mathcal{V}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$


$\llbracket \alpha \mid x : A \vdash [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow T\mathbb{N}$



$\llbracket \mid x : A \vdash \nu \alpha. [x = \alpha] : \mathbb{N} \rrbracket$

$N_A \longrightarrow T\mathbb{N}$



\mathcal{V} is a model of $\nu\rho$

\rightsquigarrow Name-abstraction:


$$\frac{\vec{\alpha}\alpha \mid \Gamma \vdash M : B}{\vec{\alpha} \mid \Gamma \vdash \nu\alpha.M : B} \quad \mapsto \quad \frac{\llbracket M \rrbracket : \llbracket \Gamma \rrbracket \rightarrow T\llbracket A \rrbracket}{\llbracket \nu\alpha.M \rrbracket = \langle \alpha \rangle \llbracket M \rrbracket : \llbracket \Gamma \rrbracket \rightarrow T\llbracket A \rrbracket}$$

\mathcal{V} is a model of $\nu\rho$

\rightsquigarrow Name-abstraction:

$$\frac{\vec{\alpha}\alpha \mid \Gamma \vdash M : B}{\vec{\alpha} \mid \Gamma \vdash \nu\alpha.M : B} \quad \mapsto \quad \frac{[[M]] : [[\Gamma]] \rightarrow T[[A]]}{[[\nu\alpha.M]] = \langle \alpha \rangle [[M]] : [[\Gamma]] \rightarrow T[[A]]}$$

$$[[\Gamma]] \xrightarrow{f} T[[A]]$$





\mathcal{V} is a model of $\nu\rho$

\rightsquigarrow Name-abstraction:

$$\frac{\vec{\alpha}\alpha \mid \Gamma \vdash M : B}{\vec{\alpha} \mid \Gamma \vdash \nu\alpha.M : B} \quad \mapsto \quad \frac{\llbracket M \rrbracket : \llbracket \Gamma \rrbracket \rightarrow T[A]}{\llbracket \nu\alpha.M \rrbracket = \langle \alpha \rangle \llbracket M \rrbracket : \llbracket \Gamma \rrbracket \rightarrow T[A]}$$

$$\llbracket \Gamma \rrbracket \xrightarrow{f} T[A]$$

$$\llbracket \Gamma \rrbracket \xrightarrow{\langle \alpha \rangle f} T[A]$$



\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$

$$\begin{array}{ccc} N_A \otimes \llbracket A \rrbracket & \xrightarrow{\text{upd}_A} & T1 \\ (\alpha, i_A)^{\vec{\alpha}} & & (\xi \Rightarrow 1 \otimes \xi) \end{array}$$

\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$

$$N_A \otimes \llbracket A \rrbracket \xrightarrow{\text{upd}_A} T1$$

$(\alpha, i_A)^{\vec{\alpha}}$
 $(\xi \Rightarrow 1 \otimes \xi)$

$*\vec{\alpha}$

\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$

$$N_A \otimes \llbracket A \rrbracket \xrightarrow{\text{upd}_A} T1$$

$(\alpha, i_A)^{\vec{\alpha}}$
 $(\xi \Rightarrow 1 \otimes \xi)$

\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$

$$N_A \otimes \llbracket A \rrbracket \xrightarrow{\text{upd}_A} T1$$

$(\alpha, i_A)^{\vec{\alpha}}$
 $(\xi \Rightarrow 1 \otimes \xi)$
 $*^{\vec{\alpha}}$
 $(*, *)^{\vec{\alpha}}$




\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$




\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$




\mathcal{V} is a model of $\nu\rho$ (2)

\rightsquigarrow Update:

$$\llbracket M \rrbracket : \Gamma \rightarrow TN_A \quad \llbracket N \rrbracket : \Gamma \rightarrow TA$$

$$\frac{}{\llbracket M := N \rrbracket : \Gamma \xrightarrow{\langle \llbracket M \rrbracket, \llbracket N \rrbracket \rangle} TN_A \otimes TA \xrightarrow{\psi} T(N_A \otimes A) \xrightarrow{T\text{upd}_A} T^2 1 \xrightarrow{\mu} T1}$$



\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:

$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$

$$N_A \xrightarrow{\text{drf}_A} T[[A]]$$

$\alpha^{\vec{\alpha}} \quad (\xi \Rightarrow [[A]] \otimes \xi)$

\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:

$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$


$$N_A \xrightarrow{\text{drf}_A} T[[A]]$$

$\alpha^{\vec{\alpha}}$ $(\xi \Rightarrow [[A]] \otimes \xi)$
 $*\vec{\alpha}$

\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:


$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$



\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:


$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$



\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:


$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$



\mathcal{V} is a model of $\nu\rho$ (3)

\rightsquigarrow Dereferencing:

$$\frac{[[M]] : [[\Gamma]] \rightarrow TN_A}{[[!M]] : [[\Gamma]] \xrightarrow{[[M]]} TN_A \xrightarrow{T\text{drf}_A} T^2[[A]] \xrightarrow{\mu} T[[A]]}$$



\mathcal{V} is a sound model

We can show *equational soundness*.

$$\llbracket M \rrbracket = \llbracket N \rrbracket \implies M \approx N$$

– *Do we also have completeness?*

\mathcal{V} is a sound model

We can show *equational soundness*.

$$\llbracket M \rrbracket = \llbracket N \rrbracket \implies M \approx N$$

– *Do we also have definability?* No.

In the reduction calculus the treatment of the store follows a specific *store-discipline*; for example,

- If a store S is updated to S' then the original store S is not accessible any more.
In strategies stores are treated as variables.

\mathcal{V} is a sound model

We can show *equational soundness*.

$$\llbracket M \rrbracket = \llbracket N \rrbracket \implies M \approx N$$

– *Do we also have definability?* No.

In the reduction calculus the treatment of the store follows a specific *store-discipline*; for example,

- If a store S is updated to S' then the original store S is not accessible any more.
In strategies stores are treated as variables.
- When the store is asked a name, it either returns its value or it deadlocks; there is no third option.
When Opponent asks the value of some name, Player is free to evade answering and play elsewhere.

Tidy strategies and Full Abstraction

We therefore restrict strategies by imposing *tidiness conditions*, obtaining thus *tidy strategies*.

Let $\mathcal{T}^{\vec{\alpha}}$ be the subcategory of $\mathcal{V}^{\vec{\alpha}}$ with objects $\llbracket A \rrbracket$ and arrows tidy strategies

Tidy strategies and Full Abstraction

We therefore restrict strategies by imposing *tidiness conditions*, obtaining thus *tidy strategies*.

Let $\mathcal{T}^{\vec{\alpha}}$ be the subcategory of $\mathcal{V}^{\vec{\alpha}}$ with objects $\llbracket A \rrbracket$ and arrows tidy strategies

$\mathcal{T} \triangleq \langle \mathcal{T}^{\vec{\alpha}}, T^{\vec{\alpha}} \rangle_{\vec{\alpha} \in \mathbb{N}^\#}$ is a sound model

If $\sigma : \llbracket A \rrbracket \rightarrow T \llbracket B \rrbracket$ is in $\mathcal{T}^{\vec{\alpha}}$ and has *finite description* then it is definable

$$\llbracket M \rrbracket \lesssim \llbracket N \rrbracket \iff M \lesssim N$$

Last slide, really

Summary:

- ↪ we introduced $\nu\rho$,
- ↪ described Nominal Games,
- ↪ constructed a store arena, and a sound model \mathcal{V} ,
- ↪ restricted to tidy strategies to obtain definability, and hence FA.

Last slide, really

Summary:

- ↪ we introduced $\nu\rho$,
- ↪ described Nominal Games,
- ↪ constructed a store arena, and a sound model \mathcal{V} ,
- ↪ restricted to tidy strategies to obtain definability, and hence FA.

Further on:

- Names for exceptions and the exception monad
- Abstract categorical models
- ‘Simpler’ quotienting to obtain the ‘nominal’ equalities
- Effectiveness and decidability

Last slide, really

Summary:

- ↪ we introduced $\nu\rho$,
- ↪ described Nominal Games,
- ↪ constructed a store arena, and a sound model \mathcal{V} ,
- ↪ restricted to tidy strategies to obtain definability, and hence FA.

THANKS

- Names for exceptions and the exception monad
- Abstract categorical models
- ‘Simpler’ quotienting to obtain the ‘nominal’ equalities
- Effectiveness and decidability

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := (\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y)); !\alpha$$

– *What does it do?*

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

$$S \vDash M \tilde{n} \tilde{m} \longrightarrow S, \beta :: V \vDash (!\beta) \tilde{n} \tilde{m}$$

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

$$\begin{aligned} S \vDash M \tilde{n} \tilde{m} &\longrightarrow S, \beta :: V \vDash (!\beta) \tilde{n} \tilde{m} \\ &\longrightarrow S, \beta :: V \vDash V \tilde{n} \tilde{m} \end{aligned}$$

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

$$S \vDash M \tilde{n} \tilde{m} \longrightarrow S, \beta :: V \vDash (!\beta) \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash V \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash (!\beta) (n \tilde{-} 1) (m \tilde{+} 1)$$

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

$$S \vDash M \tilde{n} \tilde{m} \longrightarrow S, \beta :: V \vDash (!\beta) \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash V \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash (!\beta) (n \tilde{-} 1) (m \tilde{+} 1)$$

$$\longrightarrow S, \beta :: V \vDash V (n \tilde{-} 1) (m \tilde{+} 1) \longrightarrow \dots$$

The $\nu\rho$ -calculus: An Example

$$M \triangleq \nu\alpha. \alpha := \underbrace{(\lambda x^{\mathbb{N}}, y^{\mathbb{N}}. \text{if0 } x \text{ then } y \text{ else } (!\alpha)(\text{pred } x)(\text{succ } y))}_{V}; !\alpha$$

$$\rightsquigarrow \epsilon \mid \emptyset \vdash M : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}$$

$$S \vDash M \tilde{n} \tilde{m} \longrightarrow S, \beta :: V \vDash (!\beta) \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash V \tilde{n} \tilde{m}$$

$$\longrightarrow S, \beta :: V \vDash (!\beta) (n \tilde{-} 1) (m \tilde{+} 1)$$

$$\longrightarrow S, \beta :: V \vDash V (n \tilde{-} 1) (m \tilde{+} 1) \longrightarrow \dots$$

$$\longrightarrow S, \beta :: V \vDash V \tilde{0} (m \tilde{+} n) \longrightarrow S, \beta :: V \vDash m \tilde{+} n$$