

Complexity of approximate conflict-free, linearly-ordered, and nonmonochromatic hypergraph colourings

TAMIO-VESA NAKAJIMA, University of Marburg, Germany

ZEPHYR VERWIMP, University of Oxford, United Kingdom

MARCIN WROCHNA, University of Warsaw, Poland

STANISLAV ŽIVNÝ, University of Oxford, United Kingdom

Using the algebraic approach to promise constraint satisfaction problems, we establish complexity classifications of three natural variants of hypergraph colourings: standard nonmonochromatic colourings, conflict-free colourings, and linearly-ordered colourings.

Firstly, we show that finding an ℓ -colouring of a k -colourable r -uniform hypergraph is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 3$. This provides a shorter proof of a celebrated result by Dinur, Regev, and Smyth [FOCS'02/Combinatorica'05].

Secondly, we show that finding an ℓ -conflict-free colouring of an r -uniform hypergraph that admits a k -conflict-free colouring is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 4$, except for $r = 4$ and $k = 2$ (and any ℓ); this case is solvable in polynomial time. The case of $r = 3$ is the standard nonmonochromatic colouring, and the case of $r = 2$ is the notoriously difficult open problem of approximate graph colouring.

Thirdly, we show that finding an ℓ -linearly-ordered colouring of an r -uniform hypergraph that admits a k -linearly-ordered colouring is NP-hard for all constant $3 \leq k \leq \ell$ and $r \geq 4$, thus improving on the results of Nakajima and Živný [ICALP'22/ACM ToCT'23].

CCS Concepts: • **Theory of computation** → **Design and analysis of algorithms; Problems, reductions and completeness.**

Additional Key Words and Phrases: hypergraph colourings, conflict-free colourings, unique-maximum colourings, linearly-ordered colourings

ACM Reference Format:

Tamio-Vesa Nakajima, Zephyr Verwimp, Marcin Wrochna, and Stanislav Živný. 2026. Complexity of approximate conflict-free, linearly-ordered, and nonmonochromatic hypergraph colourings. *ACM Trans. Comput. Theory* 1, 1, Article 1 (January 2026), 11 pages.

1 Introduction

Graph colouring. Graph colouring is one of the most studied computational problems: Given a graph G and an integer k , is there a k -colouring, i.e., an assignment of one of k colours to the vertices of the graph so that adjacent vertices are assigned different colours?

Deciding the existence of a 3-colouring is one of Karp's 21 NP-complete problems [19]. Since finding a graph colouring with the smallest number of colours is NP-hard, there has been much

Authors' Contact Information: Tamio-Vesa Nakajima, nakajima@uni-marburg.de, University of Marburg, Department of Mathematics and Computer Science, Marburg, Germany; Zephyr Verwimp, zephyr.verwimp@cs.ox.ac.uk, University of Oxford, Department of Computer Science, Oxford, United Kingdom; Marcin Wrochna, m.wrochna@mimuw.edu.pl, University of Warsaw, Faculty of Mathematics, Informatics and Mechanics, Warsaw, Poland; Stanislav Živný, standa.zivny@cs.ox.ac.uk, University of Oxford, Department of Computer Science, Oxford, United Kingdom.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2026 ACM.

ACM 1942-3462/2026/1-ART1

<https://doi.org/>

interest in the *approximate graph colouring* (AGC) problem: Given a graph G that admits a colouring with k colours, find a colouring with ℓ colours for some $k \leq \ell$. It is believed that for every constant $3 \leq k \leq \ell$, this problem remains NP-hard [14]. While some conditional results are known (e.g. AGC is NP-hard if we assume the *2-to-1 conjecture with perfect completeness* [10, 20], or only the *d-to-1 conjecture* [17]), proving unconditional results seems elusive. The strongest results known so far are for $\ell = 2k - 1$ [5] and $\ell = \binom{k}{\lfloor k/2 \rfloor} - 1$ [22] (the first result is stronger for $k = 3, 4$, and equal to the second for $k = 5$). As progress on proving the hardness of AGC seems to have hit a barrier, it is natural to try to attack variants of AGC, to see if any of the ideas and insights from those problems could apply to the AGC. In this paper, we will focus on hypergraph generalisations of graph colourings.

An (undirected) *r-uniform hypergraph* is a pair (V, E) , where V is the vertex set and $E \subseteq V^r$ is a set of r -tuples that is closed under permutation of coordinates. Note that a 2-uniform hypergraph is a graph. A colouring of a graph (V, E) is an assignment of colours $c(v)$ to the vertices $v \in V$ such that for every edge $(u, v) \in E$ we have $c(u) \neq c(v)$. The following three equivalent formulations describe the same condition: For every edge $(u, v) \in E$ we have that (i) the set $\{c(u), c(v)\}$ contains at least 2 elements, or (ii) some colour in the multiset $\{\{c(u), c(v)\}\}$ appears exactly once, or (iii) the largest colour in the multiset $\{\{c(u), c(v)\}\}$ appears exactly once.¹ When these three definitions are applied to hypergraphs, we get three different notions, namely nonmonochromatic (NAE) colourings, conflict-free (CF) colourings, and linearly-ordered (LO) colourings. Note that any LO colouring is a CF colouring, and any CF colouring is an NAE colouring. The three notions of colourings are different already for 4-uniform² hypergraphs.³

Promise CSPs. We will now review the most relevant literature on the three variants of hypergraph colourings. It will be convenient to present the existing results in the framework of so-called *promise constraint satisfaction problems* (PCSPs) [3, 7], as we shall use the tools developed for understanding the computational complexity of PCSPs [5]. Constraint satisfaction problems (CSPs) are problems that can be cast as homomorphisms between relational structures. We will only need a special case of relational structures that contain only one relation. Formally, a *relational structure* $\mathbf{A} = (A, R^A)$ is a pair, where A is the universe of \mathbf{A} and $R^A \subseteq A^r$ is an r -ary relation. By abuse of language, we call r the *arity* of \mathbf{A} , and we will often say \mathbf{x} is a tuple in \mathbf{A} when we mean $\mathbf{x} \in R^A$.

Note that graphs and uniform hypergraphs are relational structures, where the universe is the vertex set and the relation is the edge set – the arity of the relation is 2 for graphs and r for r -uniform hypergraphs. A *homomorphism* from one relational structure of arity r , say $\mathbf{A} = (A, R^A)$, to another of the same arity, say $\mathbf{B} = (B, R^B)$, is a map $h : A \rightarrow B$ that preserves the relation: if $(a_1, \dots, a_r) \in R^A$ then $(h(a_1), \dots, h(a_r)) \in R^B$. We denote the existence of a homomorphism from \mathbf{A} to \mathbf{B} by writing $\mathbf{A} \rightarrow \mathbf{B}$.

Given two relational structures \mathbf{A} and \mathbf{B} with $\mathbf{A} \rightarrow \mathbf{B}$, the promise constraint satisfaction problem with template (\mathbf{A}, \mathbf{B}) , denoted by $\text{PCSP}(\mathbf{A}, \mathbf{B})$, is the following computational problem. Given a relational structure \mathbf{X} with the promise $\mathbf{X} \rightarrow \mathbf{A}$, find a homomorphism from \mathbf{X} to \mathbf{B} . This is the search version of the problem. In the decision version, one is given a relational structure \mathbf{X} with the same arity as \mathbf{A} and the task is to output YES if $\mathbf{X} \rightarrow \mathbf{A}$ and No if $\mathbf{X} \not\rightarrow \mathbf{B}$. Since the decision version reduces to the search version, solving the decision version is no harder than solving the search version. All of our results will hold for both versions – hardness results will hold even for the decision version, and tractability results will hold even for the search version.

¹This notion assumes that the colours are taken from a totally ordered set, e.g., the natural numbers.

²For 3-uniform hypergraphs, NAE and CF colourings coincide, but LO colourings are different.

³Indeed, the edge (a, b, c, d) could be assigned colours $\{\{1, 1, 2, 2\}\}$ in an NAE colouring, but not in the other two; whereas the edge could be assigned colours $\{\{1, 2, 2, 2\}\}$ in a CF colouring, but not in an LO colouring.

In order to cast approximate hypergraph NAE/CF/LO-colourings as PCSPs, we will need to encode the NAE/CF/LO-colourability of a hypergraph by a homomorphism to a suitable relational structure. We will thus describe three families of relational structures capturing the three types of hypergraph colourings mentioned above (and therefore implicitly graph colouring). For any arity $r \geq 2$ and domain size k , we define:⁴

$$\begin{aligned} \text{NAE}_k^r &= (\{0, 1, \dots, k-1\}, \{(x_1, \dots, x_r) \mid \exists i, j \in [r] : x_i \neq x_j\}), \\ \text{CF}_k^r &= (\{0, 1, \dots, k-1\}, \{(x_1, \dots, x_r) \mid \exists i \in [r] \forall j \neq i \in [r] : x_i \neq x_j\}), \\ \text{LO}_k^r &= (\{0, 1, \dots, k-1\}, \{(x_1, \dots, x_r) \mid \exists i \in [r] \forall j \neq i \in [r] : x_i > x_j\}). \end{aligned}$$

Observe that an r -uniform hypergraph X has an NAE k -colouring if and only if $X \rightarrow \text{NAE}_k^r$. The analogous statement holds for CF and LO colourings. Since NAE, LO and CF colourings are all identical to graph colouring on uniformity 2, we see that k vs. ℓ AGC is the same as $\text{PCSP}(\text{NAE}_k^2, \text{NAE}_\ell^2)$ – or equivalently $\text{PCSP}(\text{CF}_k^2, \text{CF}_\ell^2)$ or $\text{PCSP}(\text{LO}_k^2, \text{LO}_\ell^2)$.

Other studied notions of hypergraph colourings include strong hypergraph colouring [6] and rainbow hypergraph colouring [2, 16, 18].

Nonmonochromatic colourings. The most studied hypergraph colourings are nonmonochromatic colourings, also known as weak hypergraph colourings. This is the weakest non-trivial restriction one can impose when colouring the vertices of a hypergraph, i.e., any type of hypergraph colouring (that excludes constant colourings) is also a nonmonochromatic colouring. As mentioned before, nonmonochromatic k -colourings of an r -uniform hypergraph correspond to homomorphisms from the hypergraph to NAE_k^r . Since nonmonochromatic colouring is NP-hard for any uniformity $r \geq 3$ and number of colours $k \geq 2$, an investigation of the approximate version led to the following result (the arity ≥ 4 case was shown in [15] without the use of topology, while the arity 3 case was shown in [11] relying on the higher chromatic number of Kneser and Schrijver graphs):

Theorem 1. $\text{PCSP}(\text{NAE}_k^r, \text{NAE}_\ell^r)$ is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 3$.

In this paper we will provide a simpler proof of this result. The proof in [11] relies on constructing a somewhat ad-hoc reduction and analysing its completeness and soundness. We recast this proof in the recent algebraic framework for PCSPs [5]. We also avoid the use of Schrijver graphs, working only with the simpler Kneser graphs, plus a (correct and very easy) case of Hedetniemi’s conjecture⁵ (cf. Lemma 9). We believe that our simplification is of interest since it replaces a more quantitative analysis of the polymorphisms (i.e. bounds are functions of the arity) with one that only deals with constant bounds everywhere.

Conflict-free colourings. A conflict-free hypergraph colouring is a colouring of the vertices in a hypergraph such that every hyperedge has at least one uniquely coloured vertex [12, 27]. As mentioned before, conflict-free k -colourings of an r -uniform hypergraph correspond to homomorphisms from the hypergraph to CF_k^r . We shall determine the complexity of $\text{PCSP}(\text{CF}_k^r, \text{CF}_\ell^r)$ for all constants $2 \leq k \leq \ell$ and $r \geq 3$ (the case of $r = 3$ corresponding to nonmonochromatic colourings, i.e., $\text{CF}_k^3 = \text{NAE}_k^3$ for every k).

After the easy observation that $\text{PCSP}(\text{CF}_k^r, \text{CF}_\ell^r)$ reduces to $\text{PCSP}(\text{CF}_k^{r+t}, \text{CF}_\ell^{r+t})$ for $t \geq 2$ (described in Lemma 6 in Section 2), Theorem 1 directly implies NP-hardness for promise conflict-free colouring for uniformity $r \geq 5$. The crux of the result is to deal with the case of uniformity $r = 4$. Note that finding a conflict-free colouring of a 4-uniform hypergraph using 2 colours is identical to solving systems of equations of the form $x + y + z + t \equiv 1 \pmod{2}$ over \mathbb{Z}_2 , and is hence in P and

⁴For any positive integer n , we write $[n]$ for the set $\{1, 2, \dots, n\}$.

⁵Note that Hedetniemi’s conjecture was proved false in the general case [26].

consequently so is $\text{PCSP}(\text{CF}_2^4, \text{CF}_\ell^4)$ for every $\ell \geq 2$. We resolve the only remaining case, showing in Section 3.3 that $\text{PCSP}(\text{CF}_k^4, \text{CF}_\ell^4)$ is NP-hard for all $3 \leq k \leq \ell$. Summarising, we have

Theorem 2. $\text{PCSP}(\text{CF}_k^r, \text{CF}_\ell^r)$ is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 3$, except for $k = 2$ and $r = 4$, which is in P.

This also immediately implies the following (much weaker) corollary, which does not appear to have been known in the CF-colouring literature.

Corollary 3. It is NP-hard to approximate the conflict-free chromatic number⁶ of a hypergraph to within any constant factor, even if it is r -uniform for some constant $r \geq 3$.

Linearly-ordered colourings. A linearly-ordered [4] (or unique-maximum [9]) hypergraph colouring is a colouring of the vertices in a hypergraph with linearly-ordered colours such that the maximum colour in every hyperedge is unique. As mentioned before, linearly-ordered k -colourings of an r -uniform hypergraph correspond to homomorphisms from the hypergraph to LO_k^r .

Barto, Battistelli, and Berg [4] conjectured that $\text{PCSP}(\text{LO}_k^3, \text{LO}_\ell^3)$ is NP-hard for all constant $2 \leq k \leq \ell$. Building on the topological methods of Krokhin, Opršal, Wrochna, and Živný [22], Filakovský, Nakajima, Opršal, Tasinato, and Wagner [13] established NP-hardness of $\text{PCSP}(\text{LO}_3^3, \text{LO}_4^3)$. This result was recently subsumed by NP-hardness of $\text{PCSP}(\text{LO}_2^3, \text{LO}_3^3)$ shown by Krokhin and Vagnozzi [21] (via a simple reduction that gives NP-hardness of $\text{PCSP}(\text{LO}_k^3, \text{LO}_{k+1}^3)$ for all $k \geq 2$).

For higher arities ($r \geq 4$), Nakajima and Živný [25] showed NP-hardness of $\text{PCSP}(\text{LO}_k^r, \text{LO}_\ell^r)$ for every $2 \leq k \leq \ell$ whenever $r \geq \ell - k + 4$. We strengthen this result, showing NP-hardness of $\text{PCSP}(\text{LO}_k^r, \text{LO}_\ell^r)$ for $3 \leq k \leq \ell$ for every $r \geq 4$.

Theorem 4. $\text{PCSP}(\text{LO}_k^r, \text{LO}_\ell^r)$ is NP-hard for all constant $3 \leq k \leq \ell$ and $r \geq 4$.

Observe that this theorem covers nearly all the cases from the result of [25]: the only case not covered is $k = 2$ and $r \geq \ell + 2$. In particular, Theorem 4 has no requirement on r in terms of ℓ , unlike the result in [25]. Indeed, Theorem 4 covers the full range of parameters except for the cases $r = 3$ or $k = 2$ (and thus the conjecture of Barto et al. remains open).

It is worth digressing somewhat to discuss the appearance of topological methods within these proofs. Our proof uses the chromatic number of the Kneser graph as an essential ingredient — this is a topological fact, and thus our proof is in some sense topological. (This is similar to the appearance of topology within hardness proofs for rainbow colourings [2].) On the other hand, the topological approach of [13], which proved that $\text{PCSP}(\text{LO}_3^3, \text{LO}_4^3)$ is NP-hard, is rather different. It assigns each relational structure an equivariant simplicial complex in a “nice enough” way so that the topological properties of these simplicial complexes imply the hardness of the original template. It would be interesting to see if these two approaches can be merged, or combined to strengthen both.

2 Preliminaries

Let (\mathbf{A}, \mathbf{B}) be a PCSP template with \mathbf{A} and \mathbf{B} of arity r . A *polymorphism* of arity $n = \text{ar}(f)$ of (\mathbf{A}, \mathbf{B}) is a function $f : A^n \rightarrow B$ such that if f is applied component-wise to any n -tuple of elements of $R^{\mathbf{A}}$ it gives an element of $R^{\mathbf{B}}$. In more detail, whenever (a_{ij}) is an $r \times n$ matrix such that every column is in $R^{\mathbf{A}}$, then f applied to the rows gives an r -tuple which is in $R^{\mathbf{B}}$. We denote by $\text{Pol}^{(n)}(\mathbf{A}, \mathbf{B})$ the collection of n -ary polymorphisms of (\mathbf{A}, \mathbf{B}) , and we let $\text{Pol}(\mathbf{A}, \mathbf{B}) = \bigcup_n \text{Pol}^{(n)}(\mathbf{A}, \mathbf{B})$.

For an n -ary function $f : A^n \rightarrow B$ and a map $\pi : [n] \rightarrow [m]$, we say that an m -ary function $g : A^m \rightarrow B$ is the *minor of f given by π* if $g(x_1, \dots, x_m) = f(x_{\pi(1)}, \dots, x_{\pi(n)})$. We write $f \xrightarrow{\pi} g$ if g is the minor of f given by π . Note that $\text{Pol}(\mathbf{A}, \mathbf{B})$ is closed under minors.

⁶That is, the minimum number of colours needed to CF-colour a given hypergraph.

We use \leq_p to denote a polynomial-time many-one reduction.

Theorem 5 ([7]). *If $\text{Pol}(\mathbf{A}, \mathbf{B}) \subseteq \text{Pol}(\mathbf{A}', \mathbf{B}')$ then $\text{PCSP}(\mathbf{A}', \mathbf{B}') \leq_p \text{PCSP}(\mathbf{A}, \mathbf{B})$.*

Lemma 6. *For any $t \geq 2$, $\text{PCSP}(\text{CF}_k^r, \text{CF}_\ell^r) \leq_p \text{PCSP}(\text{CF}_k^{r+t}, \text{CF}_\ell^{r+t})$.*

PROOF. We will use Theorem 5 and show that $\text{Pol}(\text{CF}_k^{r+t}, \text{CF}_\ell^{r+t}) \subseteq \text{Pol}(\text{CF}_k^r, \text{CF}_\ell^r)$. Suppose $f \in \text{Pol}^{(n)}(\text{CF}_k^{r+t}, \text{CF}_\ell^{r+t})$. Consider any $r \times n$ matrix A with rows $\mathbf{a}_1, \dots, \mathbf{a}_r \in [k]^n$ such that every column in the matrix has a unique entry. We can choose $\mathbf{b} \in [k]^n$ such that each column in the $(r+t) \times n$ matrix A' with rows $\mathbf{a}_1, \dots, \mathbf{a}_r$, and t copies of \mathbf{b} has a unique entry: choose element i of \mathbf{b} to be any value in $[k]$ other than the unique entry in the i -th column of A . Since f is a polymorphism of $(\text{CF}_k^r, \text{CF}_\ell^r)$, we get that $(f(\mathbf{a}_1), \dots, f(\mathbf{a}_r), f(\mathbf{b}), \dots, f(\mathbf{b}))$ has a unique entry. Since $t \geq 2$, $(f(\mathbf{a}_1), \dots, f(\mathbf{a}_r))$ must also have a unique entry. Thus $f \in \text{Pol}^{(n)}(\text{CF}_k^r, \text{CF}_\ell^r)$ as required. \square

An ℓ -chain of minors is a sequence of the form $f_0 \xrightarrow{\pi_{0,1}} f_1 \xrightarrow{\pi_{1,2}} \dots \xrightarrow{\pi_{\ell-1,\ell}} f_\ell$. We shall then write $\pi_{i,j} : [\text{ar}(f_i)] \rightarrow [\text{ar}(f_j)]$ for the composition of $\pi_{i,i+1}, \dots, \pi_{j-1,j}$, for any $0 \leq i < j \leq \ell$. Note that $f_i \xrightarrow{\pi_{i,j}} f_j$. We shall use the following NP-hardness criterion for PCSPs.

Theorem 7 ([8]). *Suppose there are constants k, ℓ and an assignment sel which, for every $f \in \text{Pol}(\mathbf{A}, \mathbf{B})$, outputs a set $\text{sel}(f) \subseteq [n]$ of size at most k , where n is the arity of f . Suppose furthermore that for every ℓ -chain of minors there is a pair $i < j$ such that $\pi_{i,j}(\text{sel}(f_i)) \cap \text{sel}(f_j) \neq \emptyset$. Then, $\text{PCSP}(\mathbf{A}, \mathbf{B})$ is NP-hard.*

For a graph G , the *chromatic number* of G , denoted by $\chi(G)$, is the smallest k such that $G \rightarrow K_k$, where K_k is the clique on k vertices. We will rely on Lovász's result for the chromatic number of Kneser graphs [23]. For $1 \leq h \leq |A|$, write $A^{(h)}$ for the family of subsets of A of size h . The *Kneser graph* is defined as $\text{KG}(A, h) = (A^{(h)}, E)$, where $\{P, Q\} \in E$ if and only if $P \cap Q = \emptyset$. For the special case $A = [n]$, we use the notation $\text{KG}(n, h) = \text{KG}([n], h)$.

Theorem 8 ([23]). $\chi(\text{KG}(n, h)) = n - 2h + 2$ for any $n \in \mathbb{N}$, $1 \leq h \leq n/2$.

Lemma 9. *Let $\chi(G) > n$. Then $\chi(G \times K_{n+1}) > n$.*⁷

PROOF. We show the contrapositive: suppose there is a homomorphism $G \times K_{n+1} \rightarrow K_n$. Equivalently, there is a homomorphism $G \rightarrow K_n^{K_{n+1}}$, where $K_n^{K_{n+1}}$ is the graph with vertex set $\{f : [n+1] \rightarrow [n]\}$, and f and g adjacent if for every distinct $i, j \in [n+1]$, $f(i) \neq g(j)$. This is equivalent since every homomorphism $f : G \times K_{n+1} \rightarrow K_n$ corresponds uniquely to the homomorphism $f' : G \rightarrow K_n^{K_{n+1}}$ given by $f'(u) = (v \mapsto f(u, v))$. There is also a homomorphism $K_n^{K_{n+1}} \rightarrow K_n$: map any $f \in V(K_n^{K_{n+1}})$ to an arbitrary repeating element in the range of f (at least one must exist by the pigeonhole principle). Thus $G \rightarrow K_n$. \square

If $\mathbf{A} \rightarrow \mathbf{A}' \rightarrow \mathbf{B}' \rightarrow \mathbf{B}$ then (\mathbf{A}, \mathbf{B}) is a *homomorphic relaxation* of $(\mathbf{A}', \mathbf{B}')$. In this case it follows from the definitions that $\text{PCSP}(\mathbf{A}, \mathbf{B}) \leq_p \text{PCSP}(\mathbf{A}', \mathbf{B}')$ [5].

3 Proofs of hardness

3.1 Avoiding sets imply hardness

Our hardness proofs will revolve around the notion of *avoiding sets* for polymorphisms [5], defined below. For $X \subseteq [n]$, we denote by $\mathbf{1}_X$ the *indicator vector* of X : it is the n -dimensional vector with $(\mathbf{1}_X)_i = 1$ for $i \in X$ and $(\mathbf{1}_X)_i = 0$ for $i \in [n] \setminus X$.

⁷Here \times denotes the tensor product of two graphs.

Definition 10. Take A so that $\{0, 1\} \subseteq A$. Let $f : A^n \rightarrow B$ and $T \subseteq B$. A T -avoiding set for f is a set $P \subseteq [n]$ such that for any $R \supseteq P$, we have $f(\mathbf{1}_R) \notin T$. For $t \in \mathbb{N}$, we call a set P t -avoiding for f if it is T -avoiding for f for some subset $T \subseteq B$ of size t .

We will first collect some simple properties of avoiding sets.

Lemma 11. Let $f : A^n \rightarrow B$ and $\ell = |B|$.

- (i) There are no ℓ -avoiding sets for f .
- (ii) $[n]$ is an $(\ell - 1)$ -avoiding set for f .
- (iii) If U is T -avoiding for f then so is every $V \supseteq U$.
- (iv) Take $\pi : [n] \rightarrow [m]$ and suppose $f \xrightarrow{\pi} g$ for some $g : A^m \rightarrow B$. Suppose $P \subseteq [n]$ is T -avoiding for f . Then $\pi(P)$ is T -avoiding for g .

PROOF. For (i), observe that any ℓ -avoiding set would imply that $f(\mathbf{1}_{[n]}) \notin B$, which is impossible. For (ii), note that $[n]$ is $(B \setminus \{f(\mathbf{1}_{[n]})\})$ -avoiding, and hence $(\ell - 1)$ -avoiding. (iii) follows from the definitions. For (iv), first observe that $g(\mathbf{1}_X) = f(\mathbf{1}_{\pi^{-1}(X)})$. For contradiction, suppose $\pi(P)$ is not T -avoiding for g , i.e., there exists $R \supseteq \pi(P)$ with $g(\mathbf{1}_R) \in T$. Then $\pi^{-1}(R) \supseteq P$, and furthermore $f(\mathbf{1}_{\pi^{-1}(R)}) = g(\mathbf{1}_R) \in T$. Hence P is not T -avoiding for f . \square

To apply Theorem 7, we want to build $\text{sel}(f)$ out of (small) avoiding sets for f . This is a good idea because avoiding sets are preserved by minors, as shown in Lemma 11 (iv). The issue is that we might have too many avoiding sets. For the polymorphisms in this paper, many (small) t -avoiding sets which are pairwise disjoint imply the existence of a (small) $(t + 1)$ -avoiding set. Thus, since there can be no sets that avoid every output in the range, as shown in Lemma 11 (i), there must be some maximal t for which a (small) avoiding set exists. By maximality, there cannot be too many disjoint t -avoiding sets. Thus, we can build $\text{sel}(f)$ out of these disjoint “maximally avoiding” sets.

Theorem 12. Let (\mathbf{A}, \mathbf{B}) be a PCSP template with $\{0, 1\} \subseteq A$ and $\ell = |B|$. Suppose that there exist constants $N, \{\alpha_t\}_{t=1}^{\ell}, \{\beta_t\}_{t=1}^{\ell}$ such that every $f \in \text{Pol}(\mathbf{A}, \mathbf{B})$ has the following properties:

- (1) f has a 1-avoiding set of size $\leq \beta_1$.
- (2) If f is of arity $\geq N$ and has a disjoint family of $> \alpha_t$ many t -avoiding sets, all of size $\leq \beta_t$, then f has a $(t + 1)$ -avoiding set of size $\leq \beta_{t+1}$.

Then, $\text{PCSP}(\mathbf{A}, \mathbf{B})$ is NP-hard.

PROOF. For each $f \in \text{Pol}(\mathbf{A}, \mathbf{B})$, define $t(f)$ to be the maximal t such that f has a t -avoiding set of size $\leq \beta_t$. By Assumption 1 and the lack of ℓ -avoiding sets (cf. Lemma 11 (i)), $t(f)$ exists and $1 \leq t(f) < \ell$. For each $f \in \text{Pol}(\mathbf{A}, \mathbf{B})$, let \mathcal{F}_f be a maximal disjoint family of $t(f)$ -avoiding sets of size $\leq \beta_{t(f)}$. Define $\text{sel}(f) = \bigcup \mathcal{F}_f$. Then by Assumption 2, $|\text{sel}(f)| \leq \max\{N, \max_{1 \leq t < \ell} \alpha_t \beta_t\} =: k$.

Using Theorem 7, it remains to show that for every ℓ -chain of minors there are i, j such that $\pi_{i,j}(\text{sel}(f_i)) \cap \text{sel}(f_j) \neq \emptyset$. Let $f_0 \xrightarrow{\pi_{0,1}} f_1 \xrightarrow{\pi_{1,2}} \dots \xrightarrow{\pi_{\ell-1,\ell}} f_\ell$ be such a chain. Since $1 \leq t(f_i) < \ell$, there are distinct i, j such that $t(f_i) = t(f_j) =: t$. By Lemma 11 (iv), for every t -avoiding set P of f_i , $\pi_{i,j}(P)$ is t -avoiding for f_j . Hence by maximality of \mathcal{F}_{f_j} , $\pi_{i,j}(P)$ intersects $\text{sel}(f_j)$. Since $\text{sel}(f_i)$ is the union of such sets, certainly $\pi_{i,j}(\text{sel}(f_i))$ intersects $\text{sel}(f_j)$. \square

The rest of this paper will show hardness of certain PCSP templates by Theorem 12.

3.2 Hardness of promise nonmonochromatic colouring

In this section we prove the advertised hardness results for NAE colouring. The most important case is $\text{PCSP}(\text{NAE}_2^3, \text{NAE}_\ell^3)$ for $\ell \geq 2$; all other cases derive from this one by either gadget reductions or homomorphic relaxations.

Lemma 13. *Let $\ell \geq 2$ and $n \in \mathbb{N}$. Any $f \in \text{Pol}^{(n)}(\text{NAE}_2^3, \text{NAE}_\ell^3)$ has a 1-avoiding set of size $\leq \ell$.*

PROOF. By Lemma 11 (ii), $[n]$ is an $(\ell - 1)$ -avoiding set and hence a 1-avoiding set. Thus if $n \leq \ell$, we are done. Otherwise assume $n > \ell$. Let $h = \lceil \frac{n-\ell}{2} \rceil$. Consider the Kneser graph $\text{KG}(n, h)$, and colour each vertex P by $f(\mathbf{1}_P)$. By Theorem 8, $\chi(\text{KG}(n, h)) = n - 2h + 2 > \ell$, so there are disjoint sets $P, Q \in [n]^{\binom{h}{2}}$ with the same colour $f(\mathbf{1}_P) = f(\mathbf{1}_Q) =: b$. Let $X := [n] \setminus (P \cup Q)$.

We claim that X is a $\{b\}$ -avoiding set and thus 1-avoiding. Since $|X| = n - 2h \leq \ell$ this completes the proof. In order to prove the claim, consider any Y such that $X \subseteq Y \subseteq [n]$; we want to show that $f(\mathbf{1}_Y) \neq b$. Construct a matrix in which each column corresponds to an element $i \in [n]$ and whose rows are $\mathbf{1}_P, \mathbf{1}_Q, \mathbf{1}_Y$. Observe that since $P \cup Q \cup Y = [n]$ (as P, Q, X is a partition of $[n]$), no column contains only 0s. Similarly since $P \cap Q \cap Y = \emptyset$ (as P and Q are disjoint), no column contains only 1s. Hence all the columns of the matrix whose rows are $\mathbf{1}_P, \mathbf{1}_Q, \mathbf{1}_Y$ are tuples of NAE_2^3 . Since f is a polymorphism of $(\text{NAE}_2^3, \text{NAE}_\ell^3)$, $(f(\mathbf{1}_P), f(\mathbf{1}_Q), f(\mathbf{1}_Y))$ must be a tuple in NAE_ℓ^3 . But $f(\mathbf{1}_P) = f(\mathbf{1}_Q) = b$, so $f(\mathbf{1}_Y) \neq b$ as required. Thus X is 1-avoiding. \square

Lemma 14. *Let $1 \leq t < \ell$ and $n \geq (\ell + 1)\ell^t + \ell + 1$. Suppose $f \in \text{Pol}^{(n)}(\text{NAE}_2^3, \text{NAE}_\ell^3)$ has $> \binom{\ell}{t} \cdot \ell$ disjoint t -avoiding sets of size $\leq \ell^t$. Then f has a $(t + 1)$ -avoiding set of size $\leq \ell^{t+1}$.*

PROOF. By assumption and the pigeonhole principle, f has $\geq \ell + 1$ disjoint sets $S_1, \dots, S_{\ell+1} \subseteq [n]$ of size $\leq \ell^t$ that avoid the same $T \subseteq \{0, 1, \dots, \ell - 1\}$ of size $|T| = t$. Let $R := [n] \setminus (S_1 \cup \dots \cup S_{\ell+1})$. Let $h = \lceil \frac{|R| - \ell}{2} \rceil$. We have $h \geq 0$ by the lower bound on n .

Consider subsets of $[n]$ which are the union of exactly one of $S_1, \dots, S_{\ell+1}$, and a subset of R of size h ; let \mathcal{S} be the collection of such subsets. We want to find two disjoint sets $P, Q \in \mathcal{S}$ such that $f(\mathbf{1}_P) = f(\mathbf{1}_Q)$. Observe that there is a bijection between \mathcal{S} and the vertex set of $K_{\ell+1} \times \text{KG}(R, h)$, given by taking vertex (i, A) of $K_{\ell+1} \times \text{KG}(R, h)$ to $S_i \cup A \in \mathcal{S}$. Furthermore, this bijection extends to an isomorphism between the graph $K_{\ell+1} \times \text{KG}(R, h)$ and the graph whose vertex set is \mathcal{S} and which considers $P, Q \in \mathcal{S}$ to be adjacent if and only if they are disjoint. By Lemma 9, $\chi(K_{\ell+1} \times \text{KG}(R, h)) > \ell$, so there must exist disjoint sets $P, Q \in \mathcal{S}$ with $f(\mathbf{1}_P) = f(\mathbf{1}_Q) =: b$.

Since $S_i \subseteq P$ for some i , we have $b \notin T$. Let $X := [n] \setminus (P \cup Q)$. Identically to the reasoning in the proof of Lemma 13, observe that for any $Y \subseteq [n]$ with $X \subseteq Y$, $f(\mathbf{1}_Y) \neq b$. Moreover, $X \subseteq Y$ implies $S_j \subseteq Y$ for $(\ell + 1) - 2 \geq 1$ different values of j (i.e. those for which $S_j \not\subseteq \{P, Q\}$), hence $f(\mathbf{1}_Y) \notin T$. Thus X is $(T \cup \{b\})$ -avoiding, so $(t + 1)$ -avoiding. Finally $|X| \leq ((\ell + 1) - 2) \cdot \ell^t + |R| - 2h \leq (\ell - 1) \cdot \ell^t + \ell \leq \ell^{t+1}$. \square

Theorem 1. *$\text{PCSP}(\text{NAE}_k^r, \text{NAE}_\ell^r)$ is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 3$.*

PROOF. $\text{PCSP}(\text{NAE}_2^3, \text{NAE}_\ell^3)$ is NP-hard for all $\ell \geq 2$ by Theorem 12, Lemma 13, and Lemma 14. To extend NP-hardness to larger uniformities $r \geq 3$, it is sufficient to observe that the map sending a 3-uniform hypergraph $I = (V, E)$ to the r -uniform hypergraph $I' = (V, E')$ where $E' = \{(x, y, z, \dots, z) \mid (x, y, z) \in E\}$ describes a reduction $\text{PCSP}(\text{NAE}_2^3, \text{NAE}_\ell^3) \leq_p \text{PCSP}(\text{NAE}_2^r, \text{NAE}_\ell^r)$. Finally, we can extend NP-hardness to any pair $k \leq \ell$ since by homomorphic relaxation we have $\text{PCSP}(\text{NAE}_2^r, \text{NAE}_\ell^r) \leq_p \text{PCSP}(\text{NAE}_k^r, \text{NAE}_\ell^r)$. \square

3.3 Hardness of promise conflict-free and linearly-ordered colouring

In this section we prove the advertised hardness results for both LO and CF colourings. For the CF colourings, by Lemma 6 it suffices to establish hardness for $r = 4$. However, our proof is the same for any $r \geq 4$ and thus we will present it that way. Since $\text{LO}_k^r \rightarrow \text{CF}_k^r$, we can then do both LO and CF colourings “in one go” by proving the hardness of $\text{PCSP}(\text{LO}_3^r, \text{CF}_\ell^r)$ for all $\ell \geq 3$ and $r \geq 4$. In the following proofs, we let $\mathbf{0}$ denote the vector whose elements are all 0, and we let 2_X denote a “scaled indicator vector”: $2_X = 2 \cdot \mathbf{1}_X$.

Lemma 15. *Let $\ell \geq 3$ and $r \geq 4$. Then any $f \in \text{Pol}^{(n)}(\text{LO}_3^r, \text{CF}_\ell^r)$ has a 1-avoiding set of size $\leq \ell$.*

PROOF. By Lemma 11 (ii), $[n]$ is an $(\ell - 1)$ -avoiding set and hence a 1-avoiding set. Thus if $n \leq \ell$, we are done. Otherwise assume $n > \ell$, and set $h = \lceil \frac{n-\ell}{2} \rceil$. Consider the Kneser graph $\text{KG}(n, h)$, and colour each vertex P by $f(2_P)$. By Theorem 8, $\chi(\text{KG}(n, h)) = n - 2h + 2 > \ell$, so there exist disjoint sets $P, Q \in [n]^{(h)}$ such that $f(2_P) = f(2_Q)$. Let $X := [n] \setminus (P \cup Q)$.

Now observe that for any $Y \subseteq [n]$ with $X \subseteq Y$, all the columns of the $r \times n$ matrix whose rows are $1_Y, 2_P, 2_Q$, and $r - 3$ many copies of $\mathbf{0}$ are tuples of LO_3^r . For $i \in P$, the unique maximum is a 2 in the 2_P row; for $i \in Q$ the unique maximum is a 2 in the 2_Q row; and for $i \in X$ the unique maximum is a 1 in the 1_Y row. Hence, since f is a polymorphism of $(\text{LO}_3^r, \text{CF}_\ell^r)$, we have that $(f(1_Y), f(2_P), f(2_Q), f(\mathbf{0}), \dots, f(\mathbf{0}))$ is a tuple in CF_ℓ^r . Since $f(2_P) = f(2_Q)$ we deduce that $f(1_Y) \neq f(\mathbf{0})$. Thus it follows that X is $\{f(\mathbf{0})\}$ -avoiding, and thus 1-avoiding. Noting that X has size $n - 2h \leq \ell$ completes the proof. \square

Lemma 16. *Let $\ell \geq 3$ and $r \geq 4$. Let $1 \leq t < \ell$, and suppose that $f \in \text{Pol}^{(n)}(\text{LO}_3^r, \text{CF}_\ell^r)$ has a disjoint family of $> \binom{\ell}{t} \ell$ many t -avoiding sets, all of size $\leq t\ell$. Then f has a $(t + 1)$ -avoiding set of size $\leq (t + 1)\ell$.*

PROOF. By assumption and the pigeonhole principle, there is a family \mathcal{F} of at least $\ell + 1$ disjoint T -avoiding sets of size $\leq t\ell$ for some $T \subseteq \{0, 1, \dots, \ell - 1\}$ of size t . Let h, P, Q, X be as in the proof of Lemma 15 — thus $P, Q \in [n]^{(h)}, X \in [n]^{(n-2h)}, f(2_P) = f(2_Q)$ and P, Q, X form a partition of $[n]$. Recall that $|X| \leq \ell$.

Now, since the sets in \mathcal{F} are disjoint, at most ℓ of the sets in \mathcal{F} intersect X . Thus, since $|\mathcal{F}| \geq \ell + 1$, there is some $Z \in \mathcal{F}$ disjoint from X . Let C be any other set in \mathcal{F} and define $C' = C \cup X$. Note that $|C'| \leq (t + 1)\ell$. Note also that $f(1_Z) \notin T$ as Z is T -avoiding, hence $|T \cup \{f(1_Z)\}| = t + 1$. We will show that C' is $(T \cup \{f(1_Z)\})$ -avoiding, proving the result.

Since C is T -avoiding and $C \subseteq C'$, C' is also T -avoiding (cf. Lemma 11 (iii)). To see that C' is $\{f(1_Z)\}$ -avoiding, note that for every $D' \subseteq [n]$ with $C' \subseteq D'$, all the columns of the $r \times n$ matrix whose rows are $1_{D'}, 2_P, 2_Q$, and $r - 3$ many copies of 1_Z are tuples of LO_3^r . For $i \in P$ (respectively $i \in Q$), the unique maximum is given by a 2 in the 2_P row (respectively the 2_Q row). Since P, Q, X form a partition of $[n]$, it remains to consider $i \in X$. Since P, Q, Z are disjoint from X , all the $2_P, 2_Q, 1_Z$ rows have a 0 in these columns. On the other hand, the (unique) $1_{D'}$ row has a 1, since $i \in X \subseteq C' \subseteq D'$. Hence we see that every column in the matrix is a tuple in LO_3^r . Since f is a polymorphism of $(\text{LO}_3^r, \text{CF}_\ell^r)$, $(f(2_P), f(2_Q), f(1_{D'}), f(1_Z), \dots, f(1_Z))$ is a tuple in CF_ℓ^r .

Hence since $f(2_P) = f(2_Q)$, we have that $f(1_{D'}) \neq f(1_Z)$. Thus C' is a $(T \cup \{f(1_Z)\})$ -avoiding, so $(t + 1)$ -avoiding set as required. \square

Theorem 17. *PCSP($\text{LO}_3^r, \text{CF}_\ell^r$) is NP-hard for all constant $\ell \geq 3$ and $r \geq 4$.*

PROOF. By Theorem 12, Lemma 15 and Lemma 16. \square

Theorem 2 and Theorem 4 follow immediately:

Theorem 2. *PCSP($\text{CF}_k^r, \text{CF}_\ell^r$) is NP-hard for all constant $2 \leq k \leq \ell$ and $r \geq 3$, except for $k = 2$ and $r = 4$, which is in P.*

PROOF. The case $r = 3$ is given by Theorem 1 as $\text{NAE}_k^3 = \text{CF}_k^3$ for every $k \geq 2$. The case $r \geq 5$ and $2 \leq k \leq \ell$ follows from Theorem 1 and Lemma 6. For $r = 4$ and $k = 2$, recall from Section 1 that CF_2^4 is identical to solving systems of mod-2 equations of the form $x + y + z + t \equiv 1 \pmod{2}$. Thus, $\text{CSP}(\text{CF}_2^4)$ and also $\text{PCSP}(\text{CF}_2^4, \text{CF}_\ell^4)$ is in P by homomorphic relaxation. For $r = 4$ and $3 \leq k \leq \ell$, the result follows by Theorem 17 and by homomorphic relaxation. \square

Theorem 4. $\text{PCSP}(\text{LO}_k^r, \text{LO}_\ell^r)$ is NP-hard for all constant $3 \leq k \leq \ell$ and $r \geq 4$.

PROOF. By Theorem 17 and by homomorphic relaxation as $\text{LO}_3^r \rightarrow \text{LO}_k^r$ and $\text{LO}_\ell^r \rightarrow \text{CF}_\ell^r$. \square

3.4 Extending hardness to other templates

In the above proofs of Lemma 15 and Lemma 16, the only required property of CF_ℓ^r is that for every tuple in the relation, if the first two entries are equal, then the remaining $r - 2$ entries in the tuple cannot all be equal. Thus, the same proof also shows a stronger result:

Definition 18. Define $\text{BNAE}_\ell^{s,r-s}$ (Block-NAE) as the relational structure with domain $\{0, 1, \dots, \ell - 1\}$, with a single r -ary relation which contains the tuples for which if any s entries are the same, then the remaining $r - s$ entries cannot all be the same. (This relation with $s = 2$ is strictly larger than the relation corresponding to CF_ℓ^r when $r \geq 6$).

Then the exact same proof from Section 3.3 (replacing every occurrence of CF_ℓ^r with $\text{BNAE}_\ell^{2,r-2}$) shows that $\text{PCSP}(\text{LO}_3^r, \text{BNAE}_\ell^{2,r-2})$ is NP-hard for all constant $\ell \geq 3, r \geq 4$.

In fact, using Kneser hypergraphs and the same proof technique, one can also show NP-hardness of $\text{PCSP}(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$ for all constant $\ell \geq 3, r \geq 4, 2 \leq s \leq r - 2$.

Definition 19. The s -uniform Kneser hypergraph is defined as $\text{KG}^{(s)}(A, h) = (A^{(h)}, E)$, where $\{P_1, \dots, P_s\} \in E$ if and only if $\{P_1, \dots, P_s\}$ is a collection of pairwise disjoint subsets of A of size h . Again, for the special case $A = [n]$, we use the notation $\text{KG}^{(s)}(n, h) = \text{KG}^{(s)}([n], h)$.

There is an analogue of Lovász's theorem for Kneser hypergraphs [1].⁸

Theorem 20 ([1]). $\chi(\text{KG}^{(s)}(n, h)) = \left\lceil \frac{n-s(h-1)}{s-1} \right\rceil$ for any $n \in \mathbb{N}, 1 \leq h \leq n/s$.

The proof of hardness then continues completely analogously (and is actually a generalisation of the proof in Section 3.3), but is included for completeness.

Lemma 21. Let $\ell \geq 3, r \geq 4$, and $2 \leq s \leq r - 2$. Then any $f \in \text{Pol}^{(n)}(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$ has a 1-avoiding set of size $\leq \ell(s - 1)$.

PROOF. By Lemma 11 (ii), $[n]$ is an $(\ell - 1)$ -avoiding set and hence a 1-avoiding set. Thus if $n \leq \ell(s - 1)$, we are done. Otherwise assume $n > \ell(s - 1)$, and set $h = \lceil \frac{n-\ell(s-1)}{s} \rceil$. Consider the Kneser graph $\text{KG}^{(s)}([n], h)$, and colour each vertex P by $f(2_P)$. By Theorem 20, $\chi(\text{KG}^{(s)}(n, h)) = \left\lceil \frac{n-s(h-1)}{s-1} \right\rceil > \ell$, so there exist disjoint sets $P_1, \dots, P_s \in [n]^{(h)}$ such that $f(2_{P_1}) = f(2_{P_2}) = \dots = f(2_{P_s})$. Let $X := [n] \setminus \bigcup_{j \in [s]} P_j$.

Now observe that for any $Y \subseteq [n]$ with $X \subseteq Y$, all the columns of the $r \times n$ matrix whose rows are $1_Y, 2_{P_1}, 2_{P_2}, \dots, 2_{P_s}$, and $r - s - 1$ many copies of $\mathbf{0}$ are tuples of LO_3^r . For $i \in P_j$, the unique maximum is a 2 in the 2_{P_j} row; and for $i \in X$ the unique maximum is a 1 in the 1_Y row. Hence, since f is a polymorphism of $(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$, we have that $(f(1_Y), f(2_{P_1}), f(2_{P_2}), \dots, f(2_{P_s}), f(\mathbf{0}), \dots, f(\mathbf{0}))$ is a tuple in $\text{BNAE}_\ell^{s,r-s}$. Since $f(2_{P_1}) = f(2_{P_2}) = \dots = f(2_{P_s})$ we deduce that $f(1_Y) \neq f(\mathbf{0})$. Thus it follows that X is $\{f(\mathbf{0})\}$ -avoiding, and thus 1-avoiding. Noting that X has size $n - sh \leq \ell(s - 1)$ completes the proof. \square

Lemma 22. Let $\ell \geq 3, r \geq 4, 2 \leq s \leq r - 2$. Let $1 \leq t < \ell$, and suppose that $f \in \text{Pol}^{(n)}(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$ has a disjoint family of $> \binom{\ell}{t}(s - 1)\ell$ many t -avoiding sets, all of size $\leq t(s - 1)\ell$. Then f has a $(t + 1)$ -avoiding set of size $\leq (t + 1)(s - 1)\ell$

⁸For an s -uniform hypergraph H , the chromatic number $\chi(H)$ of H , is the smallest k such that $H \rightarrow \text{NAE}_k^s$.

PROOF. By assumption and the pigeonhole principle, there is a family \mathcal{F} of at least $(s-1)\ell + 1$ disjoint T -avoiding sets of size $\leq t(s-1)\ell$ for some $T \subseteq \{0, 1, \dots, \ell-1\}$ of size t . Let h, P_1, \dots, P_s, X be as in the proof of Lemma 21. Recall $|X| \leq (s-1)\ell$.

Now, since the sets in \mathcal{F} are disjoint, at most $(s-1)\ell$ of the sets in \mathcal{F} intersect X . Thus, since $|\mathcal{F}| \geq (s-1)\ell + 1$, there is some $Z \in \mathcal{F}$ disjoint from X . Let C be any other set in \mathcal{F} and define $C' = C \cup X$. Note that $|C'| \leq (t+1)(s-1)\ell$. Note also that $f(\mathbf{1}_Z) \notin T$ as Z is T -avoiding, hence $|T \cup \{f(\mathbf{1}_Z)\}| = t+1$. We will show that C' is $(T \cup \{f(\mathbf{1}_Z)\})$ -avoiding, proving the result.

Since C is T -avoiding and $C \subseteq C'$, C' is also T -avoiding (cf. Lemma 11 (iii)). To see that C' is $\{f(\mathbf{1}_Z)\}$ -avoiding, note that for every $D' \subseteq [n]$ with $C' \subseteq D'$, all the columns of the $r \times n$ matrix whose rows are $\mathbf{1}_{D'}, 2_{P_1}, 2_{P_2}, \dots, 2_{P_s}$, and $r-s-1$ many copies of $\mathbf{1}_Z$ are tuples of LO_3^r . For $i \in P_j$, the unique maximum is given by a 2 in the 2_{P_j} row. Since P_1, \dots, P_s, X form a partition of $[n]$, it remains to consider $i \in X$. Since P_1, \dots, P_s, Z are disjoint from X , the $\mathbf{1}_Z$ row and all the 2_{P_j} rows have a 0 in these columns. On the other hand, the (unique) $\mathbf{1}_{D'}$ row has a 1, since $i \in X \subseteq C' \subseteq D'$. Hence we see that every column in the matrix is a tuple in LO_3^r . Since f is a polymorphism of $(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$, $(f(2_{P_1}), f(2_{P_2}), \dots, f(2_{P_s}), f(\mathbf{1}_{D'}), f(\mathbf{1}_Z), \dots, f(\mathbf{1}_Z))$ is a tuple in $\text{BNAE}_\ell^{s,r-s}$.

Hence since $f(2_{P_1}) = f(2_{P_2}) = \dots = f(2_{P_s})$, we have that $f(\mathbf{1}_{D'}) \neq f(\mathbf{1}_Z)$. Thus C' is a $(T \cup \{f(\mathbf{1}_Z)\})$ -avoiding, so $(t+1)$ -avoiding set as required. \square

Theorem 23. $\text{PCSP}(\text{LO}_3^r, \text{BNAE}_\ell^{s,r-s})$ is NP-hard for all constant $\ell \geq 3, r \geq 4$, and $2 \leq s \leq r-2$.

PROOF. By Theorem 12, Lemma 21 and Lemma 22. \square

We finish with noting that our proofs can be used to show NP-hardness of more templates if the symmetry requirement is dropped, then capturing colouring variants of directed hypergraphs.

Acknowledgments

An extended abstract of a part of this work appeared in Proceedings of ICALP (B) 2025 [24]. This work was supported by UKRI EP/X024431/1 and Clarendon Fund Scholarship. Work done while Tamio-Vesa Nakajima was at the University of Oxford. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. All data is provided in full in the results section of this paper. Finally, we thank the anonymous reviewers for their detailed feedback.

References

- [1] N. Alon, P. Frankl, and L. Lovász. 1986. The Chromatic Number of Kneser Hypergraphs. *Trans. Am. Math. Soc.* 298, 1 (1986), 359–370. jstor:2000624 doi:10.2307/2000624
- [2] Per Austrin, Amey Bhangale, and Aditya Potukuchi. 2020. Improved Inapproximability of Rainbow Coloring. In *Proc. 31st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA'20)*. 1479–1495. arXiv:1810.02784 doi:10.1137/1.9781611975994.90
- [3] Per Austrin, Venkatesan Guruswami, and Johan Håstad. 2017. $(2+\epsilon)$ -Sat Is NP-hard. *SIAM J. Comput.* 46, 5 (2017), 1554–1573. doi:10.1137/15M1006507
- [4] Libor Barto, Diego Battistelli, and Kevin M. Berg. 2021. Symmetric Promise Constraint Satisfaction Problems: Beyond the Boolean Case. In *Proc. 38th International Symposium on Theoretical Aspects of Computer Science (STACS'21) (LIPIcs, Vol. 187)*. 10:1–10:16. arXiv:2010.04623 doi:10.4230/LIPIcs.STACS.2021.10
- [5] Libor Barto, Jakub Bulín, Andrei A. Krokhin, and Jakub Opršal. 2021. Algebraic approach to promise constraint satisfaction. *J. ACM* 68, 4 (2021), 28:1–28:66. arXiv:1811.00970 doi:10.1145/3457606
- [6] Joshua Brakensiek and Venkatesan Guruswami. 2016. New Hardness Results for Graph and Hypergraph Colorings. In *Proc. 31st Conference on Computational Complexity (CCC'16) (LIPIcs, Vol. 50)*. 14:1–14:27. doi:10.4230/LIPIcs.CCC.2016.14
- [7] Joshua Brakensiek and Venkatesan Guruswami. 2021. Promise Constraint Satisfaction: Algebraic Structure and a Symmetric Boolean Dichotomy. *SIAM J. Comput.* 50, 6 (2021), 1663–1700. arXiv:1704.01937 doi:10.1137/19M128212X
- [8] Alex Brandts, Marcin Wrochna, and Stanislav Živný. 2021. The complexity of promise SAT on non-Boolean domains. *ACM Trans. Comput. Theory* 13, 4 (2021), 26:1–26:20. arXiv:1911.09065 doi:10.1145/3470867

- [9] Panagiotis Cheilaris, Balázs Keszegh, and Dömötör Pálvölgyi. 2013. Unique-Maximum and Conflict-Free Coloring for Hypergraphs and Tree Graphs. *SIAM J. Discret. Math.* 27, 4 (2013), 1775–1787. doi:10.1137/120880471
- [10] Irit Dinur, Elchanan Mossel, and Oded Regev. 2009. Conditional Hardness for Approximate Coloring. *SIAM J. Comput.* 39, 3 (2009), 843–873. doi:10.1137/07068062X
- [11] Irit Dinur, Oded Regev, and Clifford Smyth. 2005. The Hardness of 3-Uniform Hypergraph Coloring. *Comb.* 25, 5 (2005), 519–535. doi:10.1007/s00493-005-0032-4
- [12] Guy Even, Zvi Lotker, Dana Ron, and Shakhar Smorodinsky. 2003. Conflict-Free Colorings of Simple Geometric Regions with Applications to Frequency Assignment in Cellular Networks. *SIAM J. Comput.* 33, 1 (2003), 94–136. doi:10.1137/S0097539702431840
- [13] Marek Filakovský, Tamio-Vesa Nakajima, Jakub Opršal, Gianluca Tasinato, and Uli Wagner. 2025. Hardness of Linearly Ordered 4-Colouring of 3-Colourable 3-Uniform Hypergraphs. *ACM Trans. Comput. Theory* (2025). arXiv:2312.12981 doi:10.1145/3779121
- [14] M. R. Garey and D. S. Johnson. 1976. The Complexity of Near-Optimal Graph Coloring. *J. ACM* 23, 1 (1976), 43–49. doi:10.1145/321921.321926
- [15] Venkatesan Guruswami, Johan Håstad, and Madhu Sudan. 2002. Hardness of Approximate Hypergraph Coloring. *SIAM J. Comput.* 31, 6 (2002), 1663–1686. doi:10.1137/S0097539700377165
- [16] Venkatesan Guruswami and Euiwoong Lee. 2018. Strong Inapproximability Results on Balanced Rainbow-Colorable Hypergraphs. *Comb.* 38, 3 (2018), 547–599. doi:10.1007/s00493-016-3383-0
- [17] Venkatesan Guruswami and Sai Sandeep. 2020. d-To-1 Hardness of Coloring 3-Colorable Graphs with $O(1)$ Colors. In *Proc. 47th International Colloquium on Automata, Languages, and Programming (ICALP'20) (LIPIcs, Vol. 168)*. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 62:1–62:12. doi:10.4230/LIPIcs.ICALP.2020.62
- [18] Venkatesan Guruswami and Sai Sandeep. 2020. Rainbow Coloring Hardness via Low Sensitivity Polymorphisms. *SIAM J. Discret. Math.* 34, 1 (2020), 520–537. doi:10.1137/19M127731X
- [19] Richard M. Karp. 1972. Reducibility among Combinatorial Problems. In *Complexity of Computer Computations: Proceedings of a symposium on the Complexity of Computer Computations*. Springer US, 85–103. doi:10.1007/978-1-4684-2001-2_9
- [20] Subhash Khot. 2002. On the power of unique 2-prover 1-round games. In *Proc. 34th Annual ACM Symposium on Theory of Computing (STOC'02)*. ACM, 767–775. doi:10.1145/509907.510017
- [21] Andrei Krokhnin and Danny Vagnozzi. 2025. Approximating 1-in-3 SAT by linearly ordered hypergraph 3-colouring is NP-hard. arXiv:2508.14606
- [22] Andrei A. Krokhnin, Jakub Opršal, Marcin Wrochna, and Stanislav Živný. 2023. Topology and adjunction in promise constraint satisfaction. *SIAM J. Comput.* 52, 1 (2023), 37–79. arXiv:2003.11351 doi:10.1137/20M1378223
- [23] László Lovász. 1978. Kneser's Conjecture, Chromatic Number, and Homotopy. *J. Comb. Theory, Series A* 25, 3 (Nov. 1978), 319–324. doi:10.1016/0097-3165(78)90022-5
- [24] Tamio-Vesa Nakajima, Zephyr Verwimp, Marcin Wrochna, and Stanislav Živný. 2025. Complexity of Approximate Conflict-Free, Linearly-Ordered, and Nonmonochromatic Hypergraph Colourings. In *Proc. 52nd International Colloquium on Automata, Languages, and Programming (ICALP'25) (LIPIcs, Vol. 334)*. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 169:1–169:10. doi:10.4230/LIPICS.ICALP.2025.169
- [25] Tamio-Vesa Nakajima and Stanislav Živný. 2023. Linearly Ordered Colourings of Hypergraphs. *ACM Trans. Comput. Theory* 13, 3–4 (2023). Issue Article no. 12. arXiv:2204.05628 doi:10.1145/3570909
- [26] Yaroslav Shitov. 2019. Counterexamples to Hedetniemi's conjecture. *Ann. Math.* 190, 2 (Sept. 2019). doi:10.4007/annals.2019.190.2.6
- [27] Shakhar Smorodinsky. 2013. Conflict-Free Coloring and its Applications. In *Geometry – Intuitive, Discrete, and Convex: A Tribute to László Fejes Tóth*, Imre Bárány, Károly J. Böröczky, Gábor Fejes Tóth, and János Pach (Eds.). Springer Berlin Heidelberg, Berlin, Heidelberg, 331–389. arXiv:1005.3616 doi:10.1007/978-3-642-41498-5_12